

On the ω -language Expressive Power of Extended Petri Nets

A. Finkel[†], G. Geeraerts^{‡1}, J.-F. Raskin^{‡1}, L. Van Begin^{‡1,2}

[†]*L.S.V., École Normale Supérieure de Cachan – 61, av. du Président Wilson,
94235 CACHAN Cedex, France*

[‡]*D.I., Université Libre de Bruxelles – bld. du Triomphe, 1050 Bruxelles, Belgium.*

Abstract

In this paper, we study the expressive power of several monotonic extensions of Petri nets. We compare the expressive power of Petri nets, Petri nets extended with *non-blocking arcs* and Petri nets extended with *transfer arcs*, in terms of ω -languages. We show that the hierarchy of expressive powers of those models is strict. To prove these results, we propose *original techniques* that rely on well-quasi orderings and monotonicity properties.

Key words: (extended) Petri nets, ω -languages.

1 Introduction

Reactive systems are non-terminating systems that interact with an environment. Those systems are often embedded in environments which are *safety critical*, making their correctness a crucial issue.

To formally reason about the correctness of such systems, we need formal models of their behaviours. At some abstract level, the behaviour of a non-terminating reactive system within its environment can be seen as an *infinite sequence of events* (usually taken within a finite set of events). The semantics of those systems is thus a (usually infinite) set of those infinite behaviours. Sets of infinite sequences of events have been studied intensively in automata theory where infinite sequences of events are called *infinite words*, and sets of such sequences are called *omega languages* (ω -languages).

If the global system (the reactive system and its environment) has a meaningful *finite state abstraction* then there are well-studied formalisms that can be used. For example, *finite state automata* allow us to specify any *omega regular language* [18]. Furthermore, as the global system is naturally composed of several components

¹ Partially supported by the FRFC grant 2.4530.02.

² Supported by a “First Europe” grant EPH3310300R0012 of the Walloon Region.

(at least two: the reactive system and its environment), it is convenient to model the reactive system and its environment compositionally by several (at least two) automata. This is possible using simple synchronization mechanisms. In the case of finite state machines, synchronizations on *common events* allow to model naturally most of the interesting communication mechanisms between processes.

Recently, a lot of research works have tried to generalize the *computer aided verification methods* that have been proposed for finite state systems toward infinite state systems. In particular, interesting positive (decidability) results have been obtained for a class of *parametric systems*. New methods have been proposed for automatically verifying temporal properties of concurrent systems containing an *arbitrary number* (parametric number) of finite-state processes that communicate. Contrary to the finite state case, three primitives of communication have been proposed:

- in [13], German *et al.* introduce a model where an arbitrary number of processes communicate via *rendez-vous* (synchronization on common events);
- in [8,9], Emerson *et al.*, and Esparza *et al.* study the automatic analysis of models where an arbitrary number of processes communicate through *rendez-vous* and *broadcasts*. A broadcast is a *non-blocking* synchronization mechanism where the emitter sends a signal to all the possibly awaiting processes, and continue its execution without waiting (whether there are receivers or not). In [4], Delzanno uses broadcast protocols to model and verify cache coherency protocols [14];
- In the model introduced in [5,16] by Delzanno *et al.*, an arbitrary number of processes can communicate thanks to *non-blocking rendez-vous* (in addition to *rendez-vous* and *broadcasts*). In a *non-blocking rendez-vous* synchronization, the sender emits an event, and if there are automata waiting for that event, one of those automata is chosen non-deterministically and synchronizes with the sender. As for broadcast, this synchronization mechanism is *non-blocking*. This model is useful to model multi-threaded programs written in JAVA where instructions like `NotifyAll` are modeled by using broadcasts and `Notify` are modeled by using *non-blocking rendez-vous*.

In all those works, the identity of individual processes is irrelevant. Hence, we can apply to all those models the so-called *counting abstraction* [13,19] and equivalently see all those models as extended Petri nets. It has been shown in previous works that *rendez-vous* can be modeled by *Petri nets* [15], *broadcasts* can be modeled by *Petri nets extended with transfer arcs*, and *non-blocking rendez-vous* can be modeled by *Petri nets extended with non-blocking arcs* [5,19].

These two Petri nets extensions (and others like reset Petri nets, lossy Petri nets, ...) are monotonic and well-structured [16]. Those models have attracted a lot of attention recently [6,7,6] [17,11,12] [9] [5,16]. These papers study the main decidability problems for these models: even if the *general reachability problem* is undecidable, interesting subproblems, like *control state reachability* and *termination*, are decidable for all those models, and the *boundedness* problem is decidable for Petri nets, Petri nets with non-blocking arcs and transfer nets. However, the expressiveness of those formalisms have not been studied carefully presumably because the finite word languages definable in those formalisms are all equal to the recursively enumerable languages. Nevertheless, as recalled above, those formalisms are usually used to model non-terminating systems and so their expressive power should be measured in terms of definable omega languages.

There is currently no proof that the *expressive power* of Petri nets with transfer arcs or Petri nets with non-blocking arcs, measured in terms of *definable omega languages*, are strictly greater than the expressive power of Petri nets. In this paper, we solve this open problem. Our results are as follows. First (section 3), we show that all the omega-languages definable by Petri nets with non-blocking arcs can be recognized by Petri nets with transfer arcs, but that some languages which are definable by Petri nets with transfer arcs are not recognizable by Petri nets with non-blocking arcs (even if we allow τ -transitions). Second (section 4, we show that there exist omega languages that can be defined with Petri nets extended with non-blocking arcs and can not be defined with Petri nets (even if we allow τ -transitions). The separation of expressive power over *definable omega languages* is surprising as the expressive power of those two extended Petri net models equals, as mentioned above, the expressive power of Turing Machines when measured on *finite word languages* defined with the help of a finite accepting set of markings. We also study the expressiveness of Petri nets with reset arcs in section 5.

The techniques that we use to separate the expressive power of extended Petri nets on omega languages are based on properties of *well-quasi orderings* and monotonicity. They are, to the best of our knowledge, original in the context of (extended) Petri nets.

In this version of the paper, several technical proofs have been omitted owing to lack of space. A full version of this paper is available as Technical Report number 519 of the Computer Science Department of Brussels University³

Acknowledgements We would like to address many thanks to Raymond Devillers for proof-reading the paper and giving us many judicious advices.

2 Preliminaries

In this section, we introduce the preliminaries of the discussion. In 2.1, we introduce two Petri nets extensions (non-blocking arcs and transfer arcs) and define the notion of ω -language *accepted* by these models. In 2.2 we recall and prove a basic result on well-quasi orderings, which is the cornerstone of the proofs of sections 3 and 4.

2.1 Extended Petri nets

Definition 2.1 An **Extended Petri Net** (EPN for short) \mathcal{N} is a tuple $\langle \mathcal{P}, \mathcal{T}, \Sigma, \mathbf{m}_0 \rangle$, where $\mathcal{P} = \{p_1, p_2, \dots, p_n\}$ is a finite set of places, \mathcal{T} is finite set of transitions and Σ is a finite alphabet containing a special silent symbol τ . A **marking** of the places is a function $\mathbf{m} : \mathcal{P} \mapsto \mathbb{N}$. A marking can also be seen as a vector v such that $v^T = [\mathbf{m}(p_1), \mathbf{m}(p_2), \dots, \mathbf{m}(p_n)]$. $\mathbf{m}_0 : \mathcal{P} \mapsto \mathbb{N}$ is the initial marking. Each transition is of the form $\langle I, O, s, d, b, \lambda \rangle$, where I and $O : \mathcal{P} \mapsto \mathbb{N}$ are multi-sets of input and output places respectively. By convention, $O(p)$ (resp. $I(p)$) denotes the number of occurrences of p in O (resp. I). $s, d \in \mathcal{P} \cup \{\perp\}$ are the source and the destination places respectively, $b \in \mathbb{N} \cup \{+\infty\}$ is the bound and $\lambda \in \Sigma$ is the label of the transition.

Let us divide \mathcal{T} into \mathcal{T}_r and \mathcal{T}_e such that $\mathcal{T} = \mathcal{T}_r \cup \mathcal{T}_e$ and $\mathcal{T}_e \cap \mathcal{T}_r = \emptyset$. Without loss of generality, we assume that for each transition $\langle I, O, s, d, b, \lambda \rangle \in \mathcal{T}$, either $b = 0$ and $s = \perp = d$ (regular Petri transitions, grouped into \mathcal{T}_r); or $b > 0$,

³ which can be downloaded at: <http://www.ulb.ac.be/di/ssd/ggeeraer/papers/express.pdf>

$s \neq d$, $s \neq \perp$, $d \neq \perp$ (extended transitions, grouped into \mathcal{T}_e). We identify several non-disjoint classes of Extended Petri Nets, depending on \mathcal{T}_e :

- **Petri net** (PN for short): $\mathcal{T}_e = \emptyset$.
- **Petri net with non-blocking arcs** (PN+NBA): $\forall t = \langle I, O, s, d, b, \lambda \rangle \in \mathcal{T}_e : b = 1$.
- **Petri net with transfer arcs** (PN+T): $\forall t = \langle I, O, s, d, b, \lambda \rangle \in \mathcal{T}_e : b = +\infty$. ■

As usual, places are graphically depicted by circles; transitions by filled rectangles. For any transition $t = \langle I, O, s, d, b, \lambda \rangle$, we draw an arrow from any place $p \in I$ to transition t and from t to any place $p \in O$. For a PN+NBA (resp. PN+T), we draw a dotted (grey) arrow from s to t and from t to d (provided that $s, d \neq \perp$).

Definition 2.2 Given an extended Petri Net $\mathcal{N} = \langle \mathcal{P}, \mathcal{T}, \Sigma, \mathbf{m}_0 \rangle$, and a marking \mathbf{m} of \mathcal{N} , a transition $t = \langle I, O, s, d, b, \lambda \rangle$ is said to be *enabled in \mathbf{m}* (notation: $\mathbf{m} \xrightarrow{t}$) iff $\forall p \in \mathcal{P} : \mathbf{m}(p) \geq I(p)$. An enabled transition $t = \langle I, O, s, d, b, \lambda \rangle$ can *occur*, which deterministically transforms the marking \mathbf{m} into a new marking \mathbf{m}' (we denote this by $\mathbf{m} \xrightarrow{t} \mathbf{m}'$). \mathbf{m}' is computed as follows:

- (i) First compute \mathbf{m}_1 such that: $\forall p \in \mathcal{P} : \mathbf{m}_1(p) = \mathbf{m}(p) - I(p)$.
- (ii) Then compute \mathbf{m}_2 as follows. If $s = d = \perp$, then $\mathbf{m}_2 = \mathbf{m}_1$. Otherwise:

$$\mathbf{m}_2(s) = \begin{cases} 0 & \text{if } \mathbf{m}_1(s) \leq b \\ \mathbf{m}_1(s) - b & \text{otherwise} \end{cases} \quad \mathbf{m}_2(d) = \begin{cases} \mathbf{m}_2(d) + \mathbf{m}_1(s) & \text{if } \mathbf{m}_1(s) \leq b \\ \mathbf{m}_2(d) + b & \text{otherwise} \end{cases}$$

$$\forall p \in \mathcal{P} \setminus \{d, s\} : \mathbf{m}_2(p) = \mathbf{m}_1(p)$$

- (iii) Finally, compute \mathbf{m}' , such that $\forall p \in O : \mathbf{m}'(p) = \mathbf{m}_2(p) + O(p)$.

Let $\sigma = t_1 t_2 \dots t_n$ be a sequence of transition. We write $\mathbf{m} \xrightarrow{\sigma} \mathbf{m}'$ to mean that there exist $\mathbf{m}_1, \dots, \mathbf{m}_{n-1}$ such that $\mathbf{m} \xrightarrow{t_1} \mathbf{m}_1 \xrightarrow{t_2} \dots \xrightarrow{t_{n-1}} \mathbf{m}_{n-1} \xrightarrow{t_n} \mathbf{m}'$. ■

Definition 2.3 Let σ be a sequence of transitions. $\Lambda(\sigma)$ is defined inductively as follows (where λ_i denotes the label of t_i). If $\sigma = t_1$, then $\Lambda(\sigma) = \varepsilon$ if $\lambda_1 = \tau$; $\Lambda(\sigma) = \lambda_1$ otherwise. In the case where $\sigma = t_1 t_2 \dots$, then $\Lambda(\sigma) = \Lambda(t_2 \dots)$ if $\lambda_1 = \tau$; $\Lambda(\sigma) = \lambda_1 \cdot \Lambda(t_2 \dots)$ otherwise. ■

Remark that this definition is sound even in the case where σ is infinite since Λ associates one and only one word to any infinite sequence of transitions.

Definition 2.4 Let $\mathcal{N} = \langle \mathcal{P}, \mathcal{T}, \Sigma, \mathbf{m}_0 \rangle$. An infinite word x on Σ is said to be *accepted* by \mathcal{N} if there exists an infinite sequence of transitions $\sigma = t_1 t_2 \dots$ and an infinite set of markings $\{\mathbf{m}_1, \mathbf{m}_2 \dots\}$ such that $\mathbf{m}_1 \xrightarrow{t_1} \mathbf{m}_2 \xrightarrow{t_2} \mathbf{m}_3 \dots$, $\mathbf{m}_1 = \mathbf{m}_0$ and $\Lambda(\sigma) = x$. The *language* $L^\omega(\mathcal{N})$ is defined as the set of all the infinite words accepted by \mathcal{N} . The language $L_\neq^\omega(\mathcal{N})$ is the set of infinite words accepted by sequences of transitions of \mathcal{N} that *do not contain* τ -transitions. ■

By abuse of notation we also write $\mathbf{m} \xrightarrow{x} \mathbf{m}'$ to mean that there exists a *finite* sequence of transitions σ such that $\Lambda(\sigma) = x$ and $\mathbf{m} \xrightarrow{\sigma} \mathbf{m}'$, and $\mathbf{m} \xrightarrow{x'}$ to mean that we can fire the *infinite* sequence of transitions σ' (with $\Lambda(\sigma') = x'$) from \mathbf{m} .

In the following, $L^\omega(\text{PN})$ (respectively $L^\omega(\text{PN+T})$, $L^\omega(\text{PN+NBA})$) denotes the set of all the ω -languages that can be recognised by a PN (respectively PN+T, and

PN+NBA). $L_{\neq}^{\omega}(\text{PN})$, $L_{\neq}^{\omega}(\text{PN+NBA})$ and $L_{\neq}^{\omega}(\text{PN+T})$ are defined similarly in the case where we disallow τ -transitions.

In the sequel a notion of ordering on the markings will appear to be useful. Let \preceq denote the quasi ordering on markings, defined as follows: let \mathbf{m} and \mathbf{m}' be two markings on the set of places \mathcal{P} , then $\mathbf{m} \preceq \mathbf{m}'$ if $\forall p \in \mathcal{P} : \mathbf{m}(p) \leq \mathbf{m}'(p)$. We come back on important properties of \preceq in 2.2.

An important property of sequences of transitions of PN is their *constant effect* (it is well-known that the effect of such a sequence, when it is enabled, can be expressed by a vector of integers stating how many tokens are removed and put in each place). In the case of PN+NBA or PN+T, the effect is not constant anymore, since it is dependant on the marking at the time of the firing. However, the effect of a sequence of transitions with non-blocking arcs can be *bounded*, as stated by the following Lemma.

Lemma 2.5 *Let $\mathcal{N} = \langle \mathcal{P}, \mathcal{T}, \Sigma, \mathbf{m}_0 \rangle$ be a PN+NBA, and let σ be a finite sequence of transitions of \mathcal{N} that contains n occurrences of transitions in \mathcal{T}_e . Let \mathbf{m}_1 , \mathbf{m}'_1 , \mathbf{m}_2 and \mathbf{m}'_2 be four makings such that (i) $\mathbf{m}_1 \xrightarrow{\sigma} \mathbf{m}'_1$, (ii) $\mathbf{m}_2 \xrightarrow{\sigma} \mathbf{m}'_2$ and (iii) $\mathbf{m}_2 \succ \mathbf{m}_1$. Then, for every place $p \in \mathcal{P}$: $\mathbf{m}'_2(p) - \mathbf{m}'_1(p) \geq \mathbf{m}_2(p) - \mathbf{m}_1(p) - n$.*

Proof. Let us consider a place $p \in \mathcal{P}$. First, we remark that when we fire σ from \mathbf{m}_2 instead of \mathbf{m}_1 , its Petri net arcs will have the same effect on p . On the other hand, since we want to find a lower bound on $\mathbf{m}'_2(p) - \mathbf{m}'_1(p)$, we consider the situation where no non-blocking arcs affect p when σ is fired from \mathbf{m}_1 , but they all remove one token from p when σ is fired from \mathbf{m}_2 . In the latter case, the effect of σ on p is $\mathbf{m}'_1(p) - \mathbf{m}_1(p) - n$. We obtain thus: $\mathbf{m}'_2(p) \geq \max\{\mathbf{m}_2(p) + \mathbf{m}'_1(p) - \mathbf{m}_1(p) - n, 0\}$. Hence $\mathbf{m}'_2(p) \geq \mathbf{m}'_1(p) + \mathbf{m}_2(p) - \mathbf{m}_1(p) - n$, and thus: $\mathbf{m}'_2(p) - \mathbf{m}'_1(p) \geq \mathbf{m}_2(p) - \mathbf{m}_1(p) - n$. \square

2.2 Properties of infinite sequences on well-quasi ordered elements

Following [10,1], \preceq is a *well-quasi ordering* (wqo for short). This means that \preceq is a reflexive and transitive relation such that for any infinite sequence $\mathbf{m}_1, \mathbf{m}_2, \dots$ there is $i < j$ such that $\mathbf{m}_i \preceq \mathbf{m}_j$. Hence, we get this property on \preceq :

Lemma 2.6 *Given an infinite sequence of markings $\mathbf{m}_1, \mathbf{m}_2, \dots$ we can always extract an infinite sub-sequence $\mathbf{m}_{i_1}, \mathbf{m}_{i_2}, \dots$ ($\forall j : i_j < i_{j+1}$) such that for all place p , either $\mathbf{m}_{i_j}(p) < \mathbf{m}_{i_{j+1}}(p)$ for all $j \geq 1$ or $\mathbf{m}_{i_j}(p) = \mathbf{m}_{i_{j+1}}(p)$ for all $j \geq 1$.*

The following Lemma is easy to prove [16].

Lemma 2.7 (Monotonicity) *Let \mathbf{m}_1 , \mathbf{m}_2 and \mathbf{m}'_1 be three markings of an EPN, such that $\mathbf{m}_1 \preceq \mathbf{m}_2$ and $\mathbf{m}_1 \xrightarrow{t} \mathbf{m}'_1$ for some transition t of the EPN. Then, there exists \mathbf{m}'_2 such that $\mathbf{m}_2 \xrightarrow{t} \mathbf{m}'_2$ and $\mathbf{m}'_1 \preceq \mathbf{m}'_2$.*

3 PN+T are more expressive than PN+NBA

In this section, one will find the first important result of the paper (as stated by Theorem 3.6): PN+T are strictly more expressive, on ω -languages, than PN+NBA. We prove this in two steps. First, we show that any ω -language accepted by a PN+NBA can be accepted by a PN+T (this is the purpose of Lemma 3.1 and Theorem 3.2). Then, we prove the strictness of the inclusion thanks to the PN+T




Fig. 1. A PN+NBA \mathcal{N} (a) and the corresponding PN+T \mathcal{N}' (b)

\mathcal{N}_1 of Fig. 2 (a). Namely, we show that $L^\omega(\mathcal{N}_1)$ contains at least the words $(\mathbf{a}^k \mathbf{b}^k)^\omega$, for any $k \geq 1$ (Lemma 3.3). On the other hand we show that \mathcal{N}_1 rejects the words whose prefix belongs to $(\mathbf{a}^{n_3} \mathbf{b}^{n_3})^* \mathbf{a}^{n_3} (\mathbf{b}^{n_1} \mathbf{a}^{n_1})^+ \mathbf{b}^{n_2}$ with $n_1 < n_2 < n_3$ (Lemma 3.4). We finally show that any PN+NBA accepting words of the form $(\mathbf{a}^k \mathbf{b}^k)^\omega$ also has to accept words whose prefix belongs to $(\mathbf{a}^{n_3} \mathbf{b}^{n_3})^* \mathbf{a}^{n_3} (\mathbf{b}^{n_1} \mathbf{a}^{n_1})^+ \mathbf{b}^{n_2}$ with $n_1 < n_2 < n_3$. Since \mathcal{N}_1 rejects the latter, we conclude that no PN+NBA can accept $L^\omega(\mathcal{N}_1)$.

3.1 PN+NBA are not more expressive than PN+T.

Let us consider a PN+NBA $\mathcal{N} = \langle \mathcal{P}, \mathcal{T}, \Sigma, \mathbf{m}_0 \rangle$, and let us show how to transform it into a PN+T \mathcal{N}' such that $L^\omega(\mathcal{N}) = L^\omega(\mathcal{N}')$.

Let us consider the partition of \mathcal{T} into \mathcal{T}_e and \mathcal{T}_r as defined in Definition 2.1, and a new place p_{Tr} (the trash place). We now show how to build $\mathcal{N}' = \langle \mathcal{P}', \mathcal{T}', \Sigma, \mathbf{m}'_0 \rangle$. First, $\mathcal{P}' = \mathcal{P} \cup \{p_{Tr}\}$. For each transition $t = \langle I, O, s, d, 1, \lambda \rangle$ in \mathcal{T}_e , we put in \mathcal{T}' two new transitions $t_l = \langle I, O, s, p_{Tr}, +\infty, \lambda \rangle$ and $t_e = \langle I_e, O_e, \perp, \perp, +\infty, \lambda \rangle$, such that: $\forall p \in \mathcal{P} : (p \neq s \Rightarrow I_e(p) = I(p) \wedge p \neq d \Rightarrow O_e(p) = O(p))$, $I_e(s) = I(s) + 1$ and $O_e(d) = O(d) + 1$. We also add into \mathcal{T}' all the transitions of \mathcal{T}_r . Finally, $\forall p \in \mathcal{P} : \mathbf{m}'_0(p) = \mathbf{m}_0(p)$ and $\mathbf{m}'_0(p_{Tr}) = 0$. Fig. 1 illustrates the construction.

Lemma 3.1 $L^\omega(\mathcal{N}) = L^\omega(\mathcal{N}')$.

Proof. [$L^\omega(\mathcal{N}) \subseteq L^\omega(\mathcal{N}')$] We show that, for every infinite sequence of transitions σ of \mathcal{N} , we can find a sequence of transitions σ' of \mathcal{N}' such that $\Lambda(\sigma) = \Lambda(\sigma')$.

Let us define the function $f : \mathcal{T} \times \mathbb{N}^{|\mathcal{P}|} \rightarrow \mathcal{T}'$ such that $\forall t \in \mathcal{T}_r : f(t, \mathbf{m}) = t$ and $\forall t = \langle O, I, s, d, 1, \lambda \rangle \in \mathcal{T}_e : f(t, \mathbf{m}) = t_e$, if $\mathbf{m}(s) > I(s)$ (the non-blocking arc still has an effect after the firing of the Petri part of the transition); and $f(t, \mathbf{m}) = t_l$, otherwise.

Let $\sigma = \mathbf{m}_0 \xrightarrow{t_1} \mathbf{m}_1 \xrightarrow{t_2} \dots \xrightarrow{t_n} \mathbf{m}_n \xrightarrow{t_{n+1}} \dots$ be a sequence of \mathcal{N} . Then we may see that $\sigma' = \mathbf{m}'_0 \xrightarrow{f(t_1, \mathbf{m}'_0)} \mathbf{m}'_1 \xrightarrow{f(t_2, \mathbf{m}'_1)} \dots \xrightarrow{f(t_n, \mathbf{m}'_{n-1})} \mathbf{m}'_n \xrightarrow{f(t_{n+1}, \mathbf{m}'_n)} \dots$ is a sequence of \mathcal{N}' , where \mathbf{m}'_i is such that $\mathbf{m}'_i(p) = \mathbf{m}_i(p)$ for all $p \in \mathcal{P}$ and $\mathbf{m}'_i(p_{Tr}) = 0$ for all $i \geq 1$. Since we have $\forall i \geq 1 : \Lambda(t_i) = \Lambda(f(t_i, \mathbf{m}_{i-1}))$, we conclude that $\Lambda(\sigma) = \Lambda(\sigma')$, hence $L^\omega(\mathcal{N}) \subseteq L^\omega(\mathcal{N}')$.

[$L^\omega(\mathcal{N}') \subseteq L^\omega(\mathcal{N})$] We show that, for every infinite sequence of transitions σ' of \mathcal{N}' , we can find a sequence of transitions σ of \mathcal{N} such that $\Lambda(\sigma') = \Lambda(\sigma)$.

We define the function $g : \mathcal{T}' \rightarrow \mathcal{T}$ such that for all $t \in \mathcal{T}_r$: $g(t) = t$ and for all $t \in \mathcal{T}_e$: $g(t_e) = g(t_l) = t$. Moreover, we define the relation $\preceq_{\mathcal{P}}$ that compares two markings only on the places that are in \mathcal{P} . Thus, if \mathbf{m} is defined on \mathcal{P} and \mathbf{m}' on \mathcal{P} with $\mathcal{P}' \subseteq \mathcal{P}$, $\mathbf{m}' \preceq_{\mathcal{P}'}$ \mathbf{m} iff $\forall p \in \mathcal{P}' : \mathbf{m}'(p) \leq \mathbf{m}(p)$. Moreover, $\mathbf{m}' \prec_{\mathcal{P}'}$ \mathbf{m} iff $\mathbf{m}' \preceq_{\mathcal{P}'}$ \mathbf{m} but $\mathbf{m} \not\preceq_{\mathcal{P}'}$ \mathbf{m}' .

Let $\sigma' = \mathbf{m}'_0 \xrightarrow{t_1} \mathbf{m}'_1 \xrightarrow{t_2} \dots \xrightarrow{t_n} \mathbf{m}'_n \xrightarrow{t_{n+1}} \dots$ be a sequence of \mathcal{N}' . Then, there exist $\mathbf{m}_1, \mathbf{m}_2, \dots$ in \mathcal{N} such that we have $\mathbf{m}_0 \xrightarrow{g(t_1)} \mathbf{m}_1 \xrightarrow{g(t_2)} \dots \xrightarrow{g(t_n)} \mathbf{m}_n \xrightarrow{g(t_{n+1})} \dots$

We prove that this sequence exists by contradiction. Suppose that it is not the case, i.e. there exists $i \geq 0$ such that $g(t_{i+1})$ is not firable from \mathbf{m}_i . Let us show by induction on the indexes, that $\mathbf{m}'_j \preceq_{\mathcal{P}} \mathbf{m}_j$ for all j such that $0 \leq j \leq i$.

Base case: $j = 0$. The base case is trivially verified.

Induction step: $j = k$. By induction hypothesis, we have: $\forall 0 \leq j \leq k-1 : \mathbf{m}'_j \preceq_{\mathcal{P}} \mathbf{m}_j$. In the case where $t_k = \langle I, O, s, d, b, \lambda \rangle$ (from \mathbf{m}'_{k-1}) has the same effect on \mathcal{P} than $g(t_k)$ (from \mathbf{m}_{k-1}), we directly have that $\mathbf{m}'_k \preceq_{\mathcal{P}} \mathbf{m}_k$. This happens if t_k is a regular Petri transition or if $\mathbf{m}_{k-1}(s) = \mathbf{m}'_{k-1}(s) = I(s)$.

Otherwise t_k has a transfer arc and we must consider two cases:

- The transfer of t_k has no effect and the non-blocking arc of $g(t_k)$ moves one token from the source s to the target d , hence $I(s) = \mathbf{m}'_{k-1}(s) < \mathbf{m}_{k-1}(s)$. Since t_k and $g(t_k)$ have the same effect except that $g(t_k)$ removes one more token from s and adds one more token in d , and since $\mathbf{m}'_{k-1} \preceq_{\mathcal{P}} \mathbf{m}_{k-1}$ with $\mathbf{m}'_{k-1}(s) < \mathbf{m}_{k-1}(s)$, we conclude that $\mathbf{m}'_k \preceq_{\mathcal{P}} \mathbf{m}_k$.
- The transfer of t_k moves at least one token from the source s to p_{T_r} and the non-blocking arc of $g(t_k)$ moves one token from s to d . Since t_k and $g(t_k)$ have the same effect on the places in \mathcal{P} except that $g(t_k)$ adds one more token in d and t_k may remove more tokens from s , and since $\mathbf{m}'_{k-1} \preceq_{\mathcal{P}} \mathbf{m}_{k-1}$, we conclude that $\mathbf{m}'_k \preceq_{\mathcal{P}} \mathbf{m}_k$.

Thus $\mathbf{m}'_i \preceq_{\mathcal{P}} \mathbf{m}_i$. Since t_{i+1} is firable from \mathbf{m}'_i , we conclude that $g(t_{i+1})$ is firable from \mathbf{m}_i because $g(t_k)$ consumes no more tokens in any place p than t_k does. Hence the contradiction.

Thus, there exists $\mathbf{m}_1, \mathbf{m}_2 \dots$ such that we have $\mathbf{m}_0 \xrightarrow{g(t_1)} \mathbf{m}_1 \xrightarrow{g(t_2)} \dots \xrightarrow{g(t_n)} \mathbf{m}_n \xrightarrow{g(t_{n+1})} \dots$ in \mathcal{N} . Since $\Lambda(t_i) = \Lambda(g(t_i))$ for all $i \geq 1$, we conclude that $\Lambda(\sigma') = \Lambda(\sigma)$, hence $L^\omega(\mathcal{N}') \subseteq L^\omega(\mathcal{N})$. \square

Theorem 3.2 *For every ω -language L that is accepted by a PN+NBA, there exists a PN+T that accepts L .*

3.2 PN+T are more expressive than PN+NBA

Let us now prove that $L^\omega(\text{PN+NBA})$ is *strictly* included in $L^\omega(\text{PN+T})$. We consider the PN+T \mathcal{N}_1 presented in Fig.2 (a) with the initial marking $\mathbf{m}_0(p_1) = 1$ and $\mathbf{m}_0(p) = 0$ for $p \in \{p_2, p_3, p_4\}$. The two following Lemmata allow us to better understand the behaviour of \mathcal{N}_1 .

Lemma 3.3 *For any $k \geq 1$, the word $(\mathbf{a}^k \mathbf{b}^k)^\omega$ is accepted by \mathcal{N}_1 .*

Lemma 3.4 *Let n_1, n_2, n_3 and m be four natural numbers such that $0 < n_1 < n_2 < n_3$ and $m > 0$. Then, for any $k \geq 0$ the words $(\mathbf{a}^{n_3} \mathbf{b}^{n_3})^k \mathbf{a}^{n_3} (\mathbf{b}^{n_1} \mathbf{a}^{n_1})^m (\mathbf{b}^{n_2} \mathbf{a}^{n_2})^\omega$ are not accepted by \mathcal{N}_1 .*

We can now show that no PN+NBA can accept $L^\omega(\mathcal{N}_1)$. Remark that the proof technique used hereafter relies on Lemmata 2.6 and 2.7, and is somewhat similar to a *pumping Lemma*. To the best of our knowledge, it is the first time such a technique is applied in the context of Petri nets (and their extensions).

Lemma 3.5 *No PN+NBA accepts $L^\omega(\mathcal{N}_1)$.*

Proof. Let \mathcal{N} be a PN+NBA such that $L^\omega(\mathcal{N}_1) \subseteq L^\omega(\mathcal{N})$. We will show that this implies that $L^\omega(\mathcal{N}_1) \subsetneq L^\omega(\mathcal{N})$. As $L^\omega(\mathcal{N}_1) \subseteq L^\omega(\mathcal{N})$, by Lemma 3.3 we know that,




Fig. 2. (a) The PN+T \mathcal{N}_1 and (b) the PN+NBA \mathcal{N}_2 .

for all $k \geq 1$, the word $(\mathbf{a}^k \mathbf{b}^k)^\omega$ belongs to $L^\omega(\mathcal{N})$. Suppose that \mathbf{m}_{init} is the initial marking of \mathcal{N} . Thus, for all $k \geq 1$, there exists a marking $\widehat{\mathbf{m}}_k$, a finite sequence of transitions σ_k and a natural number ℓ_k such that:

$$\mathbf{m}_{init} \xrightarrow{(\mathbf{a}^k \mathbf{b}^k)^{\ell_k} \mathbf{a}^k} \widehat{\mathbf{m}}_k \xrightarrow{\Lambda(\sigma_k)} \widehat{\mathbf{m}}'_k \text{ and } \widehat{\mathbf{m}}'_k \succ \widehat{\mathbf{m}}_k \text{ and } \Lambda(\sigma_k) = (\mathbf{b}^k \mathbf{a}^k)^{n_k} \text{ with } n_k \geq 1$$

Indeed, suppose that it is not the case, we would have $\mathbf{m}_{init} \xrightarrow{\mathbf{a}^k} \mathbf{m}_1 \xrightarrow{\mathbf{b}^k \mathbf{a}^k} \mathbf{m}_2 \dots \xrightarrow{\mathbf{b}^k \mathbf{a}^k} \mathbf{m}_i \xrightarrow{\mathbf{b}^k \mathbf{a}^k} \dots$ such that there does not exist $1 \leq i < j$ with $\mathbf{m}_i \preceq \mathbf{m}_j$. But from Lemma 2.6, this never occurs.

Let us consider the infinite sequence $\widehat{\mathbf{m}}_1, \widehat{\mathbf{m}}_2, \dots, \widehat{\mathbf{m}}_i, \dots$. Following Lemma 2.6 again, we extract a sub-sequence from it $\widehat{\mathbf{m}}_{\rho(1)}, \widehat{\mathbf{m}}_{\rho(2)}, \dots, \widehat{\mathbf{m}}_{\rho(n)}, \dots$ such that: $\forall p \in \mathcal{P}$: either $\forall i \geq 1 : \widehat{\mathbf{m}}_{\rho(i)}(p) = \widehat{\mathbf{m}}_{\rho(i+1)}(p)$ or $\forall i \geq 1 : \widehat{\mathbf{m}}_{\rho(i)}(p) < \widehat{\mathbf{m}}_{\rho(i+1)}(p)$. Let us denote by \mathcal{P}' the set of places that strictly increase in that sequence.

Let n be the number of occurrences of transitions of \mathcal{T}_e in $\sigma_{\rho(1)}$ and let us consider $\widehat{\mathbf{m}}_{\rho(1)}, \widehat{\mathbf{m}}_{\rho(2)}, \widehat{\mathbf{m}}_{\rho(n+3)}$, and \mathbf{m} such that: $\widehat{\mathbf{m}}_{\rho(n+3)} \xrightarrow{\sigma_{\rho(1)}} \mathbf{m}$ (from Lemma 2.7, the sequence $\sigma_{\rho(1)}$ is firable since $\widehat{\mathbf{m}}_{\rho(1)} \preceq \widehat{\mathbf{m}}_{\rho(n+3)}$ and $\sigma_{\rho(1)}$ is firable from $\widehat{\mathbf{m}}_{\rho(1)}$). We first prove that $\mathbf{m} \succ \widehat{\mathbf{m}}_{\rho(2)}$.

We know that:

$$\widehat{\mathbf{m}}_{\rho(1)} \xrightarrow{\sigma_{\rho(1)}} \widehat{\mathbf{m}}'_{\rho(1)} \wedge \widehat{\mathbf{m}}'_{\rho(1)} \succ \widehat{\mathbf{m}}_{\rho(1)} \quad (1)$$

$$\forall p \in \mathcal{P}' : \widehat{\mathbf{m}}_{\rho(n+3)}(p) \geq \widehat{\mathbf{m}}_{\rho(2)}(p) + n + 1 \quad (2)$$

$$\forall p \in \mathcal{P} \setminus \mathcal{P}' : \widehat{\mathbf{m}}_{\rho(1)}(p) = \widehat{\mathbf{m}}_{\rho(2)}(p) = \widehat{\mathbf{m}}_{\rho(n+3)}(p) \quad (3)$$

Thus:

$$(a) \forall p \in \mathcal{P}' : \mathbf{m}(p) \geq \widehat{\mathbf{m}}'_{\rho(1)}(p) + (\widehat{\mathbf{m}}_{\rho(n+3)}(p) - \widehat{\mathbf{m}}_{\rho(1)}(p)) - n \quad \text{by Lemma 2.5}$$

$$\Rightarrow \forall p \in \mathcal{P}' : \mathbf{m}(p) \geq \widehat{\mathbf{m}}_{\rho(1)}(p) + (\widehat{\mathbf{m}}_{\rho(n+3)}(p) - \widehat{\mathbf{m}}_{\rho(1)}(p)) - n \quad \text{by (1)}$$

$$\Rightarrow \forall p \in \mathcal{P}' : \mathbf{m}(p) \geq \widehat{\mathbf{m}}_{\rho(n+3)}(p) - n$$

$$\Rightarrow \forall p \in \mathcal{P}' : \mathbf{m}(p) \geq \widehat{\mathbf{m}}_{\rho(2)}(p) + 1 \quad \text{by (2)}$$

$$\Rightarrow \forall p \in \mathcal{P}' : \mathbf{m}(p) > \widehat{\mathbf{m}}_{\rho(2)}(p)$$

(b) By monotonicity of PN+NBA, we have that $\mathbf{m} \succ \widehat{\mathbf{m}}'_{\rho(1)}$. Moreover, by (1), we have that $\widehat{\mathbf{m}}'_{\rho(1)} \succ \widehat{\mathbf{m}}_{\rho(1)}$. Hence, $\forall p \in \mathcal{P} : \mathbf{m}(p) \geq \widehat{\mathbf{m}}_{\rho(1)}(p)$. As a consequence, $\forall p \in \mathcal{P} \setminus \mathcal{P}' : \mathbf{m}(p) \geq \widehat{\mathbf{m}}_{\rho(2)}(p)$ from (3).

From (a) and (b), we obtain $\mathbf{m} \succ \widehat{\mathbf{m}}_{\rho(2)}$, hence $\sigma_{\rho(2)}$ is firable from \mathbf{m} . And so:

$$\mathbf{m}_{init} \xrightarrow{(\mathbf{a}^{\rho(3+n)} \mathbf{b}^{\rho(3+n)})^{\ell_{\rho(3+n)}} \mathbf{a}^{\rho(3+n)}} \widehat{\mathbf{m}}_{\rho(3+n)} \xrightarrow{(\mathbf{b}^{\rho(1)} \mathbf{a}^{\rho(1)})^{n_{\rho(1)}}} \mathbf{m} \xrightarrow{(\mathbf{b}^{\rho(2)} \mathbf{a}^{\rho(2)})^{n_{\rho(2)}}} \mathbf{m}'$$

Finally, let us prove that we can fire $\sigma_{\rho(2)}$ infinitely often from \mathbf{m}' . Since $\mathbf{m} \succ \widehat{\mathbf{m}}_{\rho(2)}$ and $\widehat{\mathbf{m}}_{\rho(2)} \xrightarrow{\sigma_{\rho(2)}} \widehat{\mathbf{m}}'_{\rho(2)}$, we have by monotonicity that $\mathbf{m}' \succ \widehat{\mathbf{m}}'_{\rho(2)} \succ \widehat{\mathbf{m}}_{\rho(2)}$, hence $\mathbf{m}' \xrightarrow{\sigma_{\rho(2)}} \mathbf{m}''$ for some marking $\mathbf{m}'' \succ \widehat{\mathbf{m}}'_{\rho(2)} \succ \widehat{\mathbf{m}}_{\rho(2)}$. Since we can repeat the reasoning infinitely often from \mathbf{m}'' , we conclude that $\sigma_{\rho(2)}$ can be fired infinitely often from \mathbf{m}'' and $(\mathbf{a}^{\rho(3+n)} \mathbf{b}^{\rho(3+n)})^{\ell_{\rho(3+n)}} \mathbf{a}^{\rho(3+n)} (\mathbf{b}^{\rho(1)} \mathbf{a}^{\rho(1)})^{n_{\rho(1)}} (\mathbf{b}^{\rho(2)} \mathbf{a}^{\rho(2)})^{\omega}$ is a word of $L^{\omega}(\mathcal{N})$ (with $\rho(3+n) > \rho(2) > \rho(1) > 0$ and $n_{\rho(1)} > 0$). But, following Lemma 3.4, this word is not in $L^{\omega}(\mathcal{N}_1)$. We conclude that $L^{\omega}(\mathcal{N}_1) \subsetneq L^{\omega}(\mathcal{N})$. \square

We can now state the main Theorem of this section, which stems directly from Theorem 3.2 and Lemma 3.5:

Theorem 3.6 *PN+T are more expressive, on infinite words, than PN+NBA, i.e.: $L^{\omega}(\text{PN+NBA}) \subsetneq L^{\omega}(\text{PN+T})$.*

Since our construction proposed to build a PN+T \mathcal{N}' from a PN+NBA \mathcal{N} that has the same ω -language than \mathcal{N} does not use τ -transitions, \mathcal{N}_1 contains no τ -transitions and since we have made no assumptions regarding the τ -transitions in the previous proofs, we obtain:

Corollary 3.7 *PN+T without τ -transitions are more expressive on infinite words than PN+NBA, i.e.: $L^{\omega}_{\neq}(\text{PN+NBA}) \subsetneq L^{\omega}_{\neq}(\text{PN+T})$.*

4 PN+NBA are more expressive than PN

In this section we prove that the class of ω -languages accepted by any PN+NBA strictly contains the class of ω -languages accepted by any PN.

The strategy adopted in the proof is similar to the one we have used in section 3. We look into the PN+NBA \mathcal{N}_2 of Fig. 2 (b), and prove it accepts every words of the form $\mathbf{i}^k \mathbf{s} (\mathbf{a}^k \mathbf{c} \mathbf{b}^k \mathbf{d})^{\omega}$, for $k \geq 1$ (Lemma 4.1), but rejects words of the form $\mathbf{i}^{n_3} \mathbf{s} (\mathbf{a}^{n_3} \mathbf{c} \mathbf{b}^{n_3} \mathbf{d})^m \mathbf{a}^{n_3} \mathbf{c} (\mathbf{b}^{n_1} \mathbf{d} \mathbf{a}^{n_1} \mathbf{c})^k (\mathbf{b}^{n_2} \mathbf{d} \mathbf{a}^{n_2} \mathbf{c})^{\omega}$, for k big enough, and $0 < n_1 < n_2 < n_3$ (Lemma 4.2). Then, we prove Lemma 4.3, stating that any PN accepting at least the words of the first form must also accept the words of the latter form. We conclude that no PN can accept $L^{\omega}(\mathcal{N}_2)$. Since any PN is also a PN+NBA, the inclusion is immediate, and we obtain Theorem 4.4, that states the strictness of the inclusion $L^{\omega}(\text{PN}) \subsetneq L^{\omega}(\text{PN+NBA})$.

Let us consider the PN+NBA \mathcal{N}_2 in Figure 2 (b), with the initial marking \mathbf{m}_0 such that $\mathbf{m}_0(p_1) = 1$ and $\mathbf{m}_0(p) = 0$ for $p \in \{p_2, p_3, p_4, p_5, p_6\}$.

Lemma 4.1 *For any $k \geq 0$, the word $\mathbf{i}^k \mathbf{s} (\mathbf{a}^k \mathbf{c} \mathbf{b}^k \mathbf{d})^{\omega}$ is accepted by \mathcal{N}_2 .*

Lemma 4.2 *Let n_1, n_2 and n_3 be three natural numbers such that $0 < n_1 < n_2 < n_3$. Then, for all $m > 0, \forall k \geq n_3 - n_1 - 1$: the words*

$$\mathbf{i}^{n_3} \mathbf{s} (\mathbf{a}^{n_3} \mathbf{c} \mathbf{b}^{n_3} \mathbf{d})^m \mathbf{a}^{n_3} \mathbf{c} (\mathbf{b}^{n_1} \mathbf{d} \mathbf{a}^{n_1} \mathbf{c})^k (\mathbf{b}^{n_2} \mathbf{d} \mathbf{a}^{n_2} \mathbf{c})^{\omega}$$

are not accepted by \mathcal{N}_2 .

We are now ready to prove that no PN accepts exactly the ω -language of the PN+NBA \mathcal{N}_2 .

Lemma 4.3 *No PN accepts $L^\omega(\mathcal{N}_2)$*

Proof. Let \mathcal{N} be a PN such that $L^\omega(\mathcal{N}_2) \subseteq L^\omega(\mathcal{N})$. We will show that this implies that $L^\omega(\mathcal{N}_2) \subsetneq L^\omega(\mathcal{N})$.

Suppose that \mathbf{m}_{init} is the initial marking of \mathcal{N} . Following Lemma 4.1, since $L^\omega(\mathcal{N}_2) \subseteq L^\omega(\mathcal{N})$, we have $\forall k \geq 1 : \mathbf{i}^k \mathbf{s}(\mathbf{a}^k \mathbf{c} \mathbf{b}^k \mathbf{d})^\omega \in L(\mathcal{N})$. Thus, for all $k \geq 1$, there exists a marking $\widehat{\mathbf{m}}_k$, a sequence of transitions σ_k and a natural ℓ_k such that:

$$\mathbf{m}_{init} \xrightarrow{\mathbf{i}^k \mathbf{s}(\mathbf{a}^k \mathbf{c} \mathbf{b}^k \mathbf{d})^{\ell_k} \mathbf{a}^k \mathbf{c}} \widehat{\mathbf{m}}_k \xrightarrow{\Lambda(\sigma_k)} \widehat{\mathbf{m}}'_k \text{ with } \widehat{\mathbf{m}}_k \preceq \widehat{\mathbf{m}}'_k \text{ and } \Lambda(\sigma_k) \in (\mathbf{b}^k \mathbf{d} \mathbf{a}^k \mathbf{c})^+$$

Indeed, suppose that it is not the case, we would have $\mathbf{m}_{init} \xrightarrow{\mathbf{i}^k \mathbf{s} \mathbf{a}^k \mathbf{c}} \mathbf{m}_1 \xrightarrow{\mathbf{b}^k \mathbf{d} \mathbf{a}^k \mathbf{c}} \dots \xrightarrow{\mathbf{b}^k \mathbf{d} \mathbf{a}^k \mathbf{c}} \mathbf{m}_i \xrightarrow{\mathbf{b}^k \mathbf{d} \mathbf{a}^k \mathbf{c}} \dots$ such that there do not exist $1 \leq i < j$ with $\mathbf{m}_i \preceq \mathbf{m}_j$. But, from Lemma 2.6, this never occurs.

Let us consider the sequence $\widehat{\mathbf{m}}_1, \widehat{\mathbf{m}}_2, \widehat{\mathbf{m}}_3, \dots$. Following Lemma 2.6, we extract an infinite sub-sequence $\widehat{\mathbf{m}}_{\rho(1)}, \widehat{\mathbf{m}}_{\rho(2)}, \widehat{\mathbf{m}}_{\rho(3)}, \dots$ such that $\forall p \in P$: either $\forall i \geq 1 : \widehat{\mathbf{m}}_{\rho(i)}(p) = \widehat{\mathbf{m}}_{\rho(i+1)}(p)$ or $\forall i \geq 1 : \widehat{\mathbf{m}}_{\rho(i)}(p) < \widehat{\mathbf{m}}_{\rho(i+1)}(p)$ (each place stays constant or strictly increases along the sequence).

Since $\widehat{\mathbf{m}}_{\rho(3)} \succcurlyeq \widehat{\mathbf{m}}_{\rho(1)}$ and $\sigma_{\rho(1)}$ has a non-negative and constant effect on each place, i.e. the effect is characterized by a tuple of naturals, we can fire $\sigma_{\rho(1)}$ any number of time from $\widehat{\mathbf{m}}_{\rho(3)}$, i.e. for all $k' \geq 0$ we have $\widehat{\mathbf{m}}_{\rho(3)} \xrightarrow{(\sigma_{\rho(1)})^{k'}} \mathbf{m}^{k'}$ with $\mathbf{m}^{k'} \succcurlyeq \widehat{\mathbf{m}}_{\rho(3)}$. Since $\widehat{\mathbf{m}}_{\rho(3)} \succcurlyeq \widehat{\mathbf{m}}_{\rho(2)}$ and $\sigma_{\rho(2)}$ has a constant non-negative effect on each place, $\sigma_{\rho(2)}$ can be fired infinitely often from $\mathbf{m}^{k'}$ for any $k' \geq 1$. Thus:

$$\mathbf{m}_{init} \xrightarrow{\mathbf{i}^{\rho(3)} \mathbf{s}(\mathbf{a}^{\rho(3)} \mathbf{c} \mathbf{b}^{\rho(3)} \mathbf{d})^{\ell_{\rho(3)}} \mathbf{a}^{\rho(3)} \mathbf{c}} \widehat{\mathbf{m}}_{\rho(3)} \xrightarrow{(\sigma_{\rho(1)})^{k'}} \mathbf{m}^{k'} \xrightarrow{(\sigma_{\rho(2)})^\omega}$$

Following Lemma 4.2, if we choose k' large enough (that is, $k' \geq \rho(3) - \rho(1) - 1$), the word accepted by the previous sequence is not in $L^\omega(\mathcal{N}_2)$. We conclude that $L^\omega(\mathcal{N}_2) \subsetneq L^\omega(\mathcal{N})$. \square

Theorem 4.4 *PN+NBA are more expressive, on infinite words, than PN, i.e.: $L^\omega(\text{PN}) \subsetneq L^\omega(\text{PN+NBA})$.*

Proof. As the PN class is a syntactic subclass of the PN+NBA, each PN-language is also a PN+NBA-language. On the other hand, some PN+NBA-languages are not PN-languages, by Lemma 4.3. Hence the Theorem. \square

Again, since PN is a syntactic subclass of PN+NBA and we have made no assumptions about the τ -transitions in the previous proofs, and since \mathcal{N}_2 contains no τ -transition, we obtain:

Corollary 4.5 *PN+NBA are more expressive, on infinite words and without τ -transitions than PN, i.e.: $L^\omega_{\neq}(\text{PN}) \subsetneq L^\omega_{\neq}(\text{PN+NBA})$.*

interleave with $t^s(t^c)^*t^e$. Moreover, σ' cannot have a suffix of the form $t^s(t^c)^\omega$ since t^c decreases the marking of the source place of the transfer of the corresponding transition t .

Then, we replace each σ_i of length > 1 by the transition t it corresponds to in \mathcal{N}_t . Hence, we obtain a new sequence $\sigma = t_1 t_2 \dots$ of \mathcal{N}_t . Clearly, $\Lambda(\sigma) = \Lambda(\sigma')$. Let us now prove that σ is firable, i.e. $\mathbf{m}_0 \xrightarrow{t_1} \mathbf{m}_1 \xrightarrow{t_2} \mathbf{m}_2 \xrightarrow{t_3} \dots$, by showing that $\forall i \geq 0 : \mathbf{m}'_{k_i} \preceq_{\mathcal{P}} \mathbf{m}_i$.

Base case: $i = 0$. The base case is trivially verified.

Induction Step: $i = l$. By induction hypothesis, we have that $\forall 0 \leq i \leq l-1 : \mathbf{m}'_{k_i} \preceq_{\mathcal{P}} \mathbf{m}_i$. In the case where t_l is a regular transition, it has the same effect on the places in \mathcal{P} as $\sigma_l = t'_l$ and it can occur since $\mathbf{m}'_{k_{l-1}} \preceq_{\mathcal{P}} \mathbf{m}_{l-1}$. Hence $\mathbf{m}'_{k_l} \preceq_{\mathcal{P}} \mathbf{m}_l$, by monotonicity. Otherwise t_l is an extended transition and its effect corresponds to the effect of σ_l . Let us observe the effect of σ_l : some tokens will be taken from s (the source place of the transfer) and put into d (the destination) by t_l^c . Finally, the tokens remaining in s will be removed by the reset arc of t_l^e . Hence, σ_l removes the same number of tokens from s than t_l , and cannot put more tokens in d than t_l does. Moreover, the effect of σ_l on the other places is the same than t_l . Thus $\mathbf{m}'_{k_l} \preceq_{\mathcal{P}} \mathbf{m}_l$. \square

Theorem 5.2 *PN+R are as expressive as PN+T on infinite words with τ -transitions.*

Proof. As any PN+R is a special case of PN+T, we have that $L^\omega(\text{PN+R}) \subseteq L^\omega(\text{PN+T})$. The other direction stems from Lemma 5.1. \square

In the case where we disallow τ -transitions, the previous construction doesn't allow to prove whether $L^\omega_{\neq}(\text{PN+T}) \subseteq L^\omega_{\neq}(\text{PN+R})$ or not. However, we have that $L^\omega(\text{PN+NBA}) \subsetneq L^\omega(\text{PN+R})$, since the PN+T \mathcal{N}_1 we have used in the proof of Lemma 3.5 satisfies our definition of PN+R (in this case, the place p_4 is the trashcan) and has no τ -transitions.

6 Conclusion

In the introduction of this paper, we have recalled how important extended Petri nets are to study the *non-terminating* behaviour of concurrent systems made up of an arbitrary number of communicating processes (once abstracted thanks to predicate- and counting- abstraction techniques [2]). Our aim was thus to study and classify the expressive powers of these models, as far as ω -languages are concerned. This goal has been thoroughly fulfilled. Indeed, we have proved in section 3 that any ω -language accepted by a PN+NBA can be accepted by a PN+T, but that there exist ω -languages that are recognised by a PN+T but not by a PN+NBA. A similar result has been demonstrated for PN+NBA and PN in section 4. These results hold with or without τ -transitions. Finally, in section 5 we have drawn a link between these results and the class of PN+R.

Future works In [15], Peterson studies different classes of *finite words* languages of PN and Ciardo [3] extends the study to Petri nets *with marking-dependant arc multiplicity*, which subsume the four classes we have studied here. The paper states some relations between these classes, but keeps several questions open. To the best of our knowledge, most of them are still open, and we strive for applying the new proof techniques developed in this paper to solve those open problems.

References

- [1] Abdulla, P. A., K. Cerans, B. Jonsson and Y.-K. Tsay, *General Decidability Theorems for Infinite-state Systems*, in: *Proceedings of the 11th Annual Symposium on Logic in Computer Science (LICS'96)* (1996), pp. 313–321.
- [2] Ball, T., S. Chaki and S. Rajamani, *Parameterized Verification of Multithreaded Software Libraries*, in: *Proceedings of the Fifth International Conference on Tools and Algorithms for Construction and Analysis of Systems (TACAS 2001)*, LNCS **2031** (2001), pp. 158–173.
- [3] Ciardo, G., *Petri nets with marking-dependent arc multiplicity: properties and analysis*, in: *Proceedings of the 15th International Conference on Applications and Theory of Petri Nets (ICATPN 94)*, LNCS **815** (1994), pp. 179–198.
- [4] Delzanno, G., *Automatic Verification of Parameterized Cache Coherence Protocols*, in: *Proceedings of the 12th International Conference on Computer Aided Verification (CAV 2000)*, LNCS **1855** (2000), pp. 53–68.
- [5] Delzanno, G., J.-F. Raskin and L. Van Begin, *Towards the Automated Verification of Multithreaded Java Programs*, in: *Proceedings of the International Conference on Tools and Algorithms for Construction and Analysis of Systems (TACAS 2002)*, LNCS **2280** (2002), pp. 173–187.
- [6] Dufourd, C., A. Finkel and P. Schnoebelen, *Reset Nets Between Decidability and Undecidability*, in: *In Proceedings of the 25th International Colloquium on Automata, Languages, and Programming (ICALP'98)*, LNCS **1443** (1998), pp. 103–115.
- [7] Dufourd, C., P. Jančar and P. Schnoebelen, *Boundedness of reset P/T nets*, in: *Proc. 2th Int. coll. Automat, Languages and Programming (ICALP'99)*, Prague, Czech Republic, LNCS **1644** (1999), pp. 301–310.
- [8] Emerson, E. A. and K. S. Namjoshi, *On Model Checking for Non-deterministic Infinite-state Systems*, in: *Proceedings of the 13th Annual Symposium on Logic in Computer Science (LICS '98)* (1998), pp. 70–80.
- [9] Esparza, J., A. Finkel and R. Mayr, *On the Verification of Broadcast Protocols*, in: *Proceedings of the 14th Annual Symposium on Logic in Computer Science (LICS'99)* (1999), pp. 352–359.
- [10] Finkel, A., *Reduction and Covering of Infinite Reachability Trees*, Information and Computation **89** (1990), pp. 144–179.
- [11] Finkel, A. and G. Sutre, *An algorithm constructing the semilinear post* for 2-dim Reset Transfer VASS*, in: *Proc. 25th Int. Symp. Math. Found. Comp. Sci. (MFCS'2000)*, Bratislava, Slovakia, LNCS **1893** (2000), pp. 353–362.
- [12] Finkel, A. and G. Sutre, *Decidability of reachability problems for classes of two counters automata*, in: *Proc. 1th Ann. Symp. Theoretical Aspects of Computer Science (STACS'2000)*, Lille, France, LNCS **1770** (2000), pp. 346–357.

- [13] German, S. M. and A. P. Sistla, *Reasoning about Systems with Many Processes*, Journal of ACM **39** (1992), pp. 675–735.
- [14] Handy, J., “The cache memory book,” Academic Press Professional, Inc., 1993.
- [15] Peterson, J. L., “Petri Net Theory and the Modeling of Systems,” Prentice Hall, 1981.
- [16] Raskin, J.-F. and L. Van Begin, *Petri Nets with Non-blocking Arcs are Difficult to Analyse*, in: *Proceedings of the 5th International Workshop on Verification of Infinite-state Systems (INFINITY 2003)*, ENTCS **96** (2003).
- [17] Sutre, G., “Abstraction et accélération de systèmes infinis,” Ph.D. thesis, École Normale Supérieure de Cachan (2000).
- [18] Thomas, W., “Handbook of theoretical computer science (vol. B): formal models and semantics,” MIT Press, 1990 pp. 133–191.
- [19] Van Begin, L., “Efficient Verification of Counting Abstractions for Parametric systems,” Ph.D. thesis, Université Libre de Bruxelles, Belgium (2003).