

# Calcium wave pacemakers in eggs

Rémi Dumollard<sup>1,2,\*</sup>, John Carroll<sup>2</sup>, Geneviève Dupont<sup>3</sup> and Christian Sardet<sup>1</sup>

<sup>1</sup>Bio Mar Cell, Unité de Biologie du Développement UMR 7009 CNRS/Paris VI, Observatoire, Station Zoologique, Villefranche sur Mer, 06230 France

<sup>2</sup>Department of Physiology, University College London, Gower Street, London, WC1E 6BT, UK

<sup>3</sup>Unité de Chronobiologie Théorique, Faculté des Sciences, Université Libre de Bruxelles, Brussels, Belgium

\*Author for correspondence (e-mail: r.dumollard@ucl.ac.uk)

*Journal of Cell Science* 115, 3557-3564 © 2002 The Company of Biologists Ltd  
doi:10.1242/jcs.00056

## Summary

During the past 25 years, the characterization of sperm-triggered calcium signals in eggs has progressed from the discovery of a single calcium increase at fertilization in the medaka fish to the observation of repetitive calcium waves initiated by multiple meiotic calcium wave pacemakers in the ascidian. In eggs of all animal species, sperm-triggered inositol (1,4,5)-trisphosphate [Ins(1,4,5)P<sub>3</sub>] production regulates the vast array of calcium wave patterns observed in the different species. The spatial organization of calcium waves is driven either by the intracellular distribution of the calcium release machinery or by the localized and dynamic production of calcium-releasing second messengers. In the highly polarized egg cell, cortical endoplasmic reticulum (ER)-rich clusters act as pacemaker

sites dedicated to the initiation of global calcium waves. The extensive ER network made of interconnected ER-rich domains supports calcium wave propagation throughout the egg. Fertilization triggers two types of calcium wave pacemakers depending on the species: in mice, the pacemaker site in the vegetal cortex of the egg is probably a site that has enhanced sensitivity to Ins(1,4,5)P<sub>3</sub>; in ascidians, the calcium wave pacemaker may rely on a local source of Ins(1,4,5)P<sub>3</sub> production apposed to a cluster of ER in the vegetal cortex.

Key words: Fertilization, Egg, Calcium oscillations, Calcium wave pacemaker, Ins(1,4,5)P<sub>3</sub>, Endoplasmic reticulum, Cortex


## Introduction

Egg cells from every animal and plant species studied to date elicit single or repetitive Ca<sup>2+</sup> transients in response to fertilizing sperm. As an important second messenger, Ca<sup>2+</sup> triggers a broad range of cellular reactions, including contraction, secretion and gene expression (Berridge, 1997; Bootman et al., 2001). In eukaryotic cells, a calcium wave starts with an initial increase in Ca<sup>2+</sup> concentration in a restricted region of the cell (a pacemaker site), which then propagates, leading to a global Ca<sup>2+</sup> wave (Berridge, 1997; Bootman et al., 2001; McDougall et al., 2000; Marchant and Parker, 2001). Several factors are important in determining the transition from non-propagated elementary Ca<sup>2+</sup> release events (Ca<sup>2+</sup> 'puffs') to the initiation and propagation of a global Ca<sup>2+</sup> wave. They include the magnitude and kinetics of Ca<sup>2+</sup> release during each elementary event, the Ca<sup>2+</sup> sensitivity of the Ca<sup>2+</sup> release channels, the spatial organization of release sites, Ca<sup>2+</sup> sequestration and Ca<sup>2+</sup> diffusion, as well as Ca<sup>2+</sup> buffering within the cytosol (Berridge, 1997; Marchant et al., 1999; Marchant and Parker, 2001; Bootman et al., 2001). In somatic cells such as hepatocytes, acinar cells or HeLa cells, all or some of these factors contribute to create a subcellular region of higher sensitivity to the Ca<sup>2+</sup>-releasing second messengers, which becomes the pacemaker site (Rooney et al., 1990; Lee et al., 1997; Petersen et al., 1999; Thomas et al., 1999; Ito et al., 1999). In eggs, global Ca<sup>2+</sup> waves always initiate in the cortex and then propagate through the cortex or the whole cytoplasm (reviewed in Sardet et al., 1998; Stricker, 1999; McDougall et al., 2000; Kline et al., 1999; Deguchi et al., 2000). After initiation of the Ca<sup>2+</sup> wave in the cortex,

propagation is due to the sequential activation by Ca<sup>2+</sup> of Ca<sup>2+</sup>-release channels at the front of the wave (Berridge, 1997; Bootman et al., 2001). The Ca<sup>2+</sup> wave speed, which mainly depends on the rate of passive Ca<sup>2+</sup> diffusion between Ca<sup>2+</sup> release sites, can be influenced by the spatial organization of these Ca<sup>2+</sup> release sites (Bugrim et al., 1997).

In eggs, Ca<sup>2+</sup> waves triggered by sperm entry result mainly from the release of Ca<sup>2+</sup> from intracellular stores by inositol 1,4,5 trisphosphate [Ins(1,4,5)P<sub>3</sub>]-induced Ca<sup>2+</sup> release (IICR) (reviewed in Miyazaki et al., 1993; Stricker, 1999; McDougall et al., 2000). The mechanism underlying Ins(1,4,5)P<sub>3</sub> production in eggs at the time of fertilization is still intensely debated. Similarly, the nature of the sperm factor(s) inducing Ca<sup>2+</sup> release at fertilization remains elusive, although several competing groups agree that it must be a protein, possibly a form of phospholipase C or an activator of it (see Stricker, 1999; Swann and Parrington, 1999; Parrington et al., 2000; McDougall et al., 2000; Nixon et al., 2000; Runft and Jaffe, 2000; Mehlmann et al., 2001; Jaffe et al., 2001; Carroll, 2001; Runft et al., 2002). Very recently, a mammalian sperm factor was characterized and found to be a new form of PLC [PLCζ (Saunders et al., 2002)]. In most species, it seems reasonable to assume that the entering sperm delivers a factor into the egg and that this factor generates Ins(1,4,5)P<sub>3</sub> either directly or indirectly.

The first wave, which we will refer to as the 'fertilization Ca<sup>2+</sup> wave', is generally the largest and longest-lasting wave, and, in some species, it is followed by repetitive Ca<sup>2+</sup> waves of lower amplitude and shorter duration. Ca<sup>2+</sup> wave pacemakers elicit waves for minutes (20-30 minutes in



**Fig. 1.** Cortical Ca<sup>2+</sup> wave pacemakers in the ascidian and mouse egg. (A) Sperm-triggered Ca<sup>2+</sup> waves in ascidians: the meiotic Ca<sup>2+</sup> waves, composed of a fertilization wave (F) followed by repetitive Ca<sup>2+</sup> waves, are initiated by two pacemakers (MI PM<sub>asc</sub> and MII PM<sub>asc</sub>). (B) An artificial pacemaker (artPM<sub>asc</sub>, red arrowhead) can be induced in the animal pole of the egg (a) by global UV photorelease of cgPtdIns(4,5)P<sub>2</sub>. The Ca<sup>2+</sup> waves emitted by this pacemaker are preceded by a pacemaker Ca<sup>2+</sup> rise (red asterisks). (C) The mouse egg is fertilized at metaphase II and thus possesses only a MII pacemaker (MII PM<sub>mouse</sub>). After the fertilization wave (F) starting from the point of sperm entry, repetitive calcium waves emanate from the vegetal cortex of the egg. Each calcium wave is preceded by a pacemaker calcium rise (red asterisks). (D) The drawings represent schematically the organization in the ascidian (MI, MII and artPM<sub>asc</sub>) and mouse egg (MII PM<sub>mouse</sub>) of the ER network (in red) and mitochondria (in green). Red arrows indicate the direction of the waves, whereas the postulated sites of Ins(1,4,5)P<sub>3</sub> production are symbolized as purple dots. sa: sperm aster; cp: contraction pole. (E) Postulated temporal variations of [Ins(1,4,5)P<sub>3</sub>]<sub>c</sub>, which may underlie the activity of the meiotic Ca<sup>2+</sup> wave pacemakers. Two possibilities remain for the ascidian and mouse MII pacemakers: a sustained Ins(1,4,5)P<sub>3</sub> production (pink trace) or oscillatory Ins(1,4,5)P<sub>3</sub> production (purple trace).

ascidians, 45-60 minutes in some molluscs, 90 minutes in nemerteans) or hours (4 hours in mammals), and they stop operating at the end of the meiotic cell cycles (except in mammals, in which they stop several hours after completion of meiosis, at the time of pronuclei formation). The pacemaker site can be fixed in the cortex or undergo dramatic movements as the cortex is reorganized in preparation for development (Sardet et al., 2002). Ca<sup>2+</sup> wave pacemakers are either located in a region of enhanced sensitivity to Ins(1,4,5)P<sub>3</sub> or reside in the vicinity of a local source of Ins(1,4,5)P<sub>3</sub>. Here, we briefly examine how the subcellular organization of the Ca<sup>2+</sup> release machinery may create stable Ca<sup>2+</sup> wave pacemakers in the egg. We also discuss how spatially and temporally regulated production of Ins(1,4,5)P<sub>3</sub> can give rise to multiple calcium wave pacemakers in a single egg cell.

#### Different calcium wave pacemakers in different species

Two main types of egg can be distinguished with regards to their patterns of sperm-triggered Ca<sup>2+</sup> signals. Eggs of sea

urchins, amphibians, cnidarians, nematode and fish display a single Ca<sup>2+</sup> increase upon fertilization. Conversely, eggs of nemerteans, some molluscs, annelids, ascidians and mammals display repetitive Ca<sup>2+</sup> waves. In most of these eggs, the oscillations following the fertilization Ca<sup>2+</sup> wave all emanate from cortical sites distinct from the initial sperm entry site (Eckberg and Miller, 1995; Kline et al., 1999; Deguchi et al., 2000) (reviewed in Sardet, 1998; Stricker, 1999). Perhaps the most elaborate of these examples is the egg of ascidians (urochordates at the base of the vertebrate line). The mature ascidian egg is arrested in metaphase I before fertilization, and sperm entry induces two series of Ca<sup>2+</sup> waves, driving, successively, the completion of meiosis I and meiosis II (Fig. 1). The first series originates from the mobile meiosis-I-associated Ca<sup>2+</sup> wave pacemaker [the MI pacemaker (McDougall and Sardet, 1995)]. The second meiotic cycle is entrained by a second pacemaker (the MII pacemaker) stably located in the vegetal cortex (Speksnijder, 1990b; McDougall and Sardet, 1995; Dumollard and Sardet, 2001) (Figs 1 and 2).

We do not completely understand why some eggs display repetitive Ca<sup>2+</sup> waves whereas others exhibit only a single

wave. Recent work on ascidian and mouse eggs reveals that arresting the egg in meiotic metaphase is both sufficient and necessary to sustain sperm-triggered  $\text{Ca}^{2+}$  oscillations (for details, see Jones, 1998; Nixon et al., 2000; Carroll, 2001). Meiotic 'M-phase' thus favors repetitive  $\text{Ca}^{2+}$  waves (as in nemerteans, some molluscs, annelids, ascidians and mammals). By contrast, only a single large fertilization  $\text{Ca}^{2+}$  wave is observed when fertilization causes a rapid transition to an interphasic cytoplasm (<20 minutes, as in fish or amphibians) and when fertilization takes place during interphase (as in cnidarians or sea urchins). It remains to be seen whether eggs of fish or amphibians can be made to undergo  $\text{Ca}^{2+}$  oscillations when blocked in meiotic M-phase after fertilization. The data relating an M-phase stage of the cell cycle to the ability to generate multiple  $\text{Ca}^{2+}$  transients is compelling in ascidian and mouse eggs (reviewed in Nixon et al., 2000; Carroll, 2001). In these eggs, regulation of the  $\text{Ca}^{2+}$  release machinery by cell cycle factors probably participates in determining the temporal pattern of the fertilization  $\text{Ca}^{2+}$  signals. However, whether such regulation proves to be universal requires further research.

### The calcium signalling hardware in eggs

The major organelles contributing to the regulation of intracellular  $\text{Ca}^{2+}$  levels are the endoplasmic reticulum (ER), the plasma membrane and mitochondria. Eggs also possess large numbers of specific vesicular organelles (yolk platelets, pigmented vesicles and cortical granules) that contain  $\text{Ca}^{2+}$  (Gillot et al., 1991); however their role in  $\text{Ca}^{2+}$  homeostasis is unknown.

The egg cortex and cytoplasm are filled with an extensive and continuous ER network (Speksnijder et al., 1993; Jaffe and Terasaki, 1993; Terasaki et al., 1996; Terasaki et al., 2001; Stricker et al., 1998; Kline et al., 1999). The ER network contains the intracellular  $\text{Ca}^{2+}$  channels – the  $\text{Ins}(1,4,5)\text{P}_3$  receptors (IP3Rs) and ryanodine receptors (RyRs) – as well as the sarco-endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPases (SERCAs) that pump calcium back into the ER.

Among the three known isoforms of IP3R found in somatic cells (Taylor et al., 1999), IP3R1 is the most prevalent and functionally important isoform in the egg [*Xenopus* (Runft et al., 1999); mammals (Miyazaki et al., 1993; Fissore et al., 1999; Brind et al., 2000); ascidian (Kyojuka et al., 1998)]. Low levels of IP3R2 and IP3R3 have been reported in mouse eggs (Fissore et al., 1999), but their physiological roles remain unclear (Brind et al., 2000; Jellerette et al., 2000). The other family of  $\text{Ca}^{2+}$  release channels (RyRs) is present on the cortical ER of sea urchin eggs (McPherson et al., 1992) and in ascidian (Albrieux et al., 2000) and mouse eggs (Ayabe et al., 1995). However, except for sea urchins, the involvement of RyRs in the initiation and propagation of sperm-triggered calcium waves appears to be minor, and their role remains unclear (reviewed in McDougall et al., 2000).

Eggs from all animal phyla seem principally to use  $\text{Ca}^{2+}$  release from internal stores to generate single or repetitive  $\text{Ca}^{2+}$  waves at fertilization. In some species, external  $\text{Ca}^{2+}$  is also used for the fertilization wave [in molluscs (Deguchi et al., 1996) (reviewed in Sardet et al., 1998)] or contributes to the maintenance of the repetitive  $\text{Ca}^{2+}$  waves [in mice (McGuinness et al., 1996) (reviewed in Stricker, 1999)]. In many species,

voltage-operated  $\text{Ca}^{2+}$  channels [VOCC (Arnoult and Villaz, 1994; Leclerc et al., 2000)] and  $\text{Ca}^{2+}$ -release-activated  $\text{Ca}^{2+}$  (CRAC) channels that mediate so-called 'capacitative  $\text{Ca}^{2+}$  entry' (Arnoult et al., 1996; Jaconi et al., 1997; Csutora et al., 1999; Machaca et al., 2000; Putney et al., 2001) are also present, but their role at fertilization is still ill defined.

In the mature mouse egg, the physiological  $\text{Ca}^{2+}$  load is primarily cleared via SERCAs and plasma membrane  $\text{Ca}^{2+}$  ATPases (PMCAs). A minor contribution may also be provided by the plasma membrane  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (Carroll, 2000). PMCAs are probably responsible for  $\text{Ca}^{2+}$  efflux from ascidian eggs after each  $\text{Ca}^{2+}$  wave (Kuthreiber et al., 1993) as well as for the loss of total  $\text{Ca}^{2+}$  content after fertilization in sea urchin eggs (Gillot et al., 1991).


In the past few years, mitochondria have been shown to be major regulators of  $\text{Ca}^{2+}$  signals (reviewed in Rutter and Rizzuto, 2000; Rizzuto et al., 2000; Duchen, 2000). Sequestration of  $\text{Ca}^{2+}$  by mitochondria has two regulatory effects on IICR, suppressing positive and negative  $\text{Ca}^{2+}$  feedback on the opening of the IP3R. In addition, ATP production by mitochondria might provide a further means of modulating  $\text{Ca}^{2+}$  signals:  $\text{ATP}^{4-}$  sensitizes the IP3R (Mak et al., 1999; Mak et al., 2001), whereas  $\text{Mg}^{2+}$ -complexed ATP is consumed to refill the ER  $\text{Ca}^{2+}$  stores. Mitochondria can thus provide negative or positive feedback on  $\text{Ins}(1,4,5)\text{P}_3$ -mediated  $\text{Ca}^{2+}$  signals. Such negative feedback has been reported in a wide range of somatic cells. For example, initiation of global  $\text{Ca}^{2+}$  waves in myocytes preferentially occurs in mitochondrion-poor regions of the cell (Boitier et al., 1999). A positive feedback effect of mitochondria on  $\text{Ins}(1,4,5)\text{P}_3$ -mediated signals has been reported only in oligodendrocytes, in which  $\text{Ca}^{2+}$  wave initiation and amplification sites are found in mitochondrion-rich regions of the cell (Simpson et al., 1997).

Except in sea urchins, in which mitochondria are a sink for cytosolic  $\text{Ca}^{2+}$  (Eisen and Reynolds, 1985; Girard et al., 1991), the role mitochondria play in  $\text{Ca}^{2+}$  signalling in eggs remains largely obscure. In mouse eggs, collapsing mitochondrial potential impairs  $\text{Ca}^{2+}$  clearance from the cytosol (Liu et al., 2001), but no picture of the regulation of  $\text{Ca}^{2+}$  oscillations by mitochondria can be drawn from only this study. In ascidian eggs, mitochondria contribute to the activity of the second  $\text{Ca}^{2+}$  wave pacemaker both by buffering cytosolic  $\text{Ca}^{2+}$  and by locally providing ATP (R.D., unpublished). Nevertheless, an understanding of the role of mitochondria in regulating  $\text{Ca}^{2+}$  wave pacemakers will require measurement of the local intracellular  $\text{Ca}^{2+}$  concentration and the local mitochondrial ATP production in the vicinity of the IP3Rs. The recent development and subcellular targeting of GFP-based  $\text{Ca}^{2+}$  and  $\text{Ins}(1,4,5)\text{P}_3$  indicators as well as luciferase-based ATP indicators should allow the direct measurement of mitochondrial  $\text{Ca}^{2+}$  levels, intracellular ATP concentration and the  $\text{Ins}(1,4,5)\text{P}_3$  concentration in the living zygote (Hirose et al., 1999; Rutter and Rizzuto, 2000).

### $\text{Ca}^{2+}$ wave pacemakers in eggs reside in cortical ER-rich domains

Given the central role played by  $\text{Ca}^{2+}$  release from intracellular stores, the organization of the ER in eggs has received much attention. In eggs that display repetitive  $\text{Ca}^{2+}$  waves, the

**Fig. 2.**  $\text{Ca}^{2+}$  wave pacemakers in eggs. (A) A sequence showing a  $\text{Ca}^{2+}$  wave initiated by the meiosis II pacemaker (MII PM) in the vegetal pole (v) of a mouse egg injected with sperm extracts. The second polar body is visible in the animal pole (a). (B) A sequence showing two examples of fertilized ascidian eggs displaying  $\text{Ca}^{2+}$  waves initiated by the MII calcium wave pacemaker located in the vegetal contraction pole (MII PM). (C) Two examples of a UV flash releasing  $\text{cgPtdIns}(4,5)P_2$  applied between two waves, the fertilized ascidian eggs respond by eliciting a wave from the animal pole of the egg (artPM). (D) The distribution of ER and mitochondria in the contraction pole (cp). A layer of cortical ER in the cortical most layer can be seen (in red) juxtaposed to the mitochondria-rich sub-cortical region (in green). (E) Schematic representation of the contraction pole showing the microvillated plasma membrane, microfilaments (in blue), as well as the ER-rich domains in the cortex (red) and the mitochondria-rich subcortical domain (green). (F) Calcium Green/Texas Red ratiometric image of  $[\text{Ca}^{2+}]_c$ , showing the initiation of a  $\text{Ca}^{2+}$  wave elicited by the MII pacemaker in the contraction pole.



interconnected network of ER sheets and tubes is organized into ER-rich domains (also called ER clusters). The cytoplasm also has ER-poor domains, which contain high densities of mitochondria and/or other vesicular organelles (Speksnijder et al., 1993; Stricker et al., 1998; Kline et al., 1999; Dumollard and Sardet, 2001). The ER clusters are made of densely packed tubes and sheets of ER membrane (Speksnijder et al., 1993; Fissore et al., 1999; Terasaki et al., 2001; Dumollard and Sardet, 2001). In oscillating eggs [nemertean (Stricker et al., 1998); mouse (Kline et al., 1999); ascidian (Dumollard and Sardet, 2001; Sardet et al., 2002)] as well as in *Xenopus* eggs (Terasaki et al., 2001), ER-rich domains are concentrated in the 2–6  $\mu\text{m}$  thick layer beneath the plasma membrane and are more

dispersed in the deeper cytoplasm (deeper than 5  $\mu\text{m}$ ). In ascidian eggs, meiosis II  $\text{Ca}^{2+}$  waves initiate in a large cortical disc of concentrated ER tubes and sheets (20  $\mu\text{m}$  in diameter and 2–5  $\mu\text{m}$  thick) located in the vegetal contraction pole (Fig. 2B–F) (Speksnijder, 1992; McDougall and Sardet, 1995; Dumollard and Sardet, 2001). In mouse and nemertean eggs, ER clusters also line the vegetal cortex of the egg where meiotic  $\text{Ca}^{2+}$  waves are initiated (Stricker et al., 1998; Kline et al., 1999).

In *Xenopus* and mouse eggs, at least, these ER-rich domains are rich in IP3R1s (Terasaki et al., 2001; Mehlmann et al., 1996; Fissore et al., 1999). The appearance of cortical ER-rich domains during maturation correlates with an increase in

sensitivity to  $\text{Ins}(1,4,5)P_3$  and to sperm-induced  $\text{Ca}^{2+}$  release (Chiba et al., 1990; Shiraishi et al., 1995; Mehlmann and Kline, 1994; Terasaki et al., 2001) (reviewed in Sardet et al., 2002). Local injections of  $\text{Ins}(1,4,5)P_3$  and sperm extracts in mouse eggs have revealed that the egg cortex is a region of higher sensitivity to  $\text{Ins}(1,4,5)P_3$  and to sperm extracts (Oda et al., 1999). Indeed, although the abundance of ER in the egg cortex renders this region more sensitive to  $\text{Ins}(1,4,5)P_3$ , it is also exposed to the highest concentrations of  $\text{Ins}(1,4,5)P_3$  as it is closest to the source of  $\text{PtdIns}(4,5)P_2$  in the plasma membrane (Halet et al., 2002; Sardet et al., 2002).

### The organization of the ER network may regulate the $\text{Ca}^{2+}$ wave pacemakers

In several somatic cells, the location of the  $\text{Ca}^{2+}$  wave pacemakers corresponds to the area of the cell that is most sensitive to  $\text{Ins}(1,4,5)P_3$  (Ito et al., 1999; Thomas et al., 1999; Petersen et al., 1999), and  $\text{Ins}(1,4,5)P_3$ , which diffuses rapidly in the cytoplasm, is thought to act as a global messenger (Albritton et al., 1992; Kasai and Petersen, 1994).

The mouse MII pacemaker appears to be an example of this type of pacemaker. In the mature mouse egg, ER-rich domains are larger in the vegetal cortex (Kline et al., 1999), whereas mitochondria are more abundant in the animal hemisphere (Calarco, 1995; Van Blerkom et al., 2002). The MII pacemaker of the mouse egg resides in the ER-enriched vegetal cortex, which is probably a site of enhanced sensitivity to  $\text{Ins}(1,4,5)P_3$ . Therefore, similarly to the somatic cell  $\text{Ca}^{2+}$  wave pacemakers, the mouse MII pacemaker site appears to be determined by the organization of the  $\text{Ca}^{2+}$  stores of the egg, with  $\text{Ins}(1,4,5)P_3$  acting as global messenger (Fig. 1, Fig. 2A).

Interestingly, an artificial pacemaker can be induced in the ascidian egg by global uncaging of caged  $\text{Ins}(1,4,5)P_3$  ( $\text{cIns}(1,4,5)P_3$ ) or its poorly metabolised analogue  $\text{cgPtdIns}(4,5)P_2$  [caged 1-( $\alpha$ -Glycerophosphoryl)-D-myoinositol 4,5-bisphosphate,  $\text{P4}(5)$ ]. This artificial pacemaker, localized in the animal pole of the ascidian egg, functions under globally elevated  $\text{Ins}(1,4,5)P_3$  levels and thus resides in the region of highest sensitivity to  $\text{Ins}(1,4,5)P_3$  (Dumollard and Sardet, 2001) (Figs 1 and 2). In common with the mouse MII pacemaker, the location of this artificial pacemaker can be explained by asymmetries in the distribution of the ER along the animal-vegetal axis of the ascidian egg. In these eggs, the ER-rich domains invade the whole egg except for the vegetal subcortex, where most mitochondria accumulate (Fig. 2) (Dumollard and Sardet, 2001). The corollary of this is that the sperm-triggered MII pacemaker in the ascidian egg, located in the vegetal pole (the site opposite the artificial pacemaker), is not at a site of enhanced  $\text{Ins}(1,4,5)P_3$  sensitivity. This indicates that the general organization of the ER stores in these eggs is not sufficient to determine the pacemaker site. The pacemakers in the ascidian egg may then rely on mechanisms other than a global increase in  $\text{Ins}(1,4,5)P_3$  levels.

### Local and dynamic production of $\text{Ins}(1,4,5)P_3$ defines the $\text{Ca}^{2+}$ wave pacemaker site

In ascidians, the MI pacemaker stimulated by sperm entry resides in a cortical ER-rich domain that forms rapidly around the sperm nucleus and centrosome and moves with them

towards the vegetal pole (Fig. 1) (Dumollard and Sardet, 2001). This suggests that the MI pacemaker of the ascidian relies on a localised moving source of  $\text{Ins}(1,4,5)P_3$ . The ascidian MII  $\text{Ca}^{2+}$  wave pacemaker does not reside in a region of enhanced sensitivity to  $\text{Ins}(1,4,5)P_3$  (Fig. 2) (Dumollard and Sardet, 2001); it might thus require local production of  $\text{Ins}(1,4,5)P_3$  in the vegetal contraction pole. The contraction pole possesses numerous microvilli and is thus rich in  $\text{PtdIns}(4,5)P_2$  (Figs 1 and 2) (Sardet et al., 2002). Therefore, the ascidian MII pacemaker may be different from the characterized pacemakers of somatic cells and the mouse MII pacemaker, since it would rely on the apposition of cortical ER-rich clusters to a local source of  $\text{Ins}(1,4,5)P_3$  (Fig. 1) (Dumollard and Sardet, 2001).

Hypothesizing local production of  $\text{Ins}(1,4,5)P_3$  even in the large egg cell raises the question of how such a gradient is maintained. Indeed, as  $\text{Ins}(1,4,5)P_3$  diffuses rapidly through the cytosol, locally produced  $\text{Ins}(1,4,5)P_3$  would quickly invade the whole cell, making  $\text{Ins}(1,4,5)P_3$  gradients energy consuming to maintain without dynamic  $\text{Ins}(1,4,5)P_3$  production. Theoretically, repetitive  $\text{Ca}^{2+}$  waves can result from either a sustained increase in  $\text{Ins}(1,4,5)P_3$  levels or an oscillating production of  $\text{Ins}(1,4,5)P_3$  (Jacob, 1990). In ascidians, a single, large and sustained  $\text{Ins}(1,4,5)P_3$  increase (achieved by uncaging  $\text{cgPtdIns}(4,5)P_2$  in the whole egg, Fig. 1) mimics the first series of  $\text{Ca}^{2+}$  oscillations, indicating that the ascidian MI pacemaker is driven by a continuous moving source of  $\text{Ins}(1,4,5)P_3$  induced by sperm entry (Dumollard and Sardet, 2001). Similarly in mouse eggs, a slow and continuous uncaging of  $\text{cIns}(1,4,5)P_3$  can reproduce the sperm-triggered  $\text{Ca}^{2+}$  oscillations (Jones and Nixon, 2000). Therefore, the mouse MII  $\text{Ca}^{2+}$  wave pacemaker can also be regulated by a single and sustained increase in  $\text{Ins}(1,4,5)P_3$  levels (Fig. 1). Furthermore, the  $\text{Ca}^{2+}$  transients triggered by the ascidian MI and artificial pacemakers, as well as those triggered by the mouse MII pacemaker, are all preceded by a characteristic slow rise in  $\text{Ca}^{2+}$  levels called a 'pacemaker  $\text{Ca}^{2+}$  rise' (Fig. 1) (Jones and Nixon, 2000; Dumollard and Sardet, 2001). This 'pacemaker  $\text{Ca}^{2+}$  rise' is a hallmark of low-frequency (period  $>20$  seconds)  $\text{Ca}^{2+}$  oscillations generated under constantly elevated  $\text{Ins}(1,4,5)P_3$  levels (Jacob, 1990; Marchant and Parker, 2001), which further suggests that a single and sustained  $\text{Ins}(1,4,5)P_3$  increase regulates the ascidian MI pacemaker and the mouse MII pacemaker.

By contrast, the ascidian MII pacemaker cannot be reproduced by a long-lasting increase in  $\text{Ins}(1,4,5)P_3$  levels in the egg, and no 'pacemaker  $\text{Ca}^{2+}$  rise' precedes these  $\text{Ca}^{2+}$  transients (Fig. 1) (Dumollard and Sardet, 2001). An oscillating production of  $\text{Ins}(1,4,5)P_3$  from the contraction pole might underlie the activity of the ascidian  $\text{Ca}^{2+}$  wave MII pacemaker (Fig. 1). The recent finding that the mammalian sperm factor is possibly a  $\text{Ca}^{2+}$ -activated phospholipase C (PLC) (Rice et al., 2000; Saunders et al., 2002) argues in favor of  $\text{Ins}(1,4,5)P_3$  oscillations driving sperm-triggered  $\text{Ca}^{2+}$  oscillations in eggs. Indeed, the prolonged stimulation of a  $\text{Ca}^{2+}$ -activated PLC can result in  $\text{Ca}^{2+}$  oscillations regulated by an oscillating production of  $\text{Ins}(1,4,5)P_3$  (for details, see Meyer and Stryer, 1988). In addition,  $\text{Ins}(1,4,5)P_3$  oscillations regulating repetitive  $\text{Ca}^{2+}$  waves during prolonged exposure to agonists have now been observed in several types of somatic cells (Hirose et al., 1999; Nash et al., 2001). The issue of  $\text{Ins}(1,4,5)P_3$  oscillations is an intensively debated topic in cell

physiology, and the  $\text{Ca}^{2+}$  wave pacemakers in eggs could provide an invaluable experimental system to resolve such questions in the future.

### A role for vegetal $\text{Ca}^{2+}$ wave pacemakers in development

Even though the ascidian and the mouse  $\text{Ca}^{2+}$  wave pacemakers seem to rely on different mechanisms, they are both located in the vegetal cortex of the egg (as is the pacemaker in eggs of the primitive nemertean). This suggests that the location the  $\text{Ca}^{2+}$  wave pacemaker may have developmental significance.

The polarized nature of the calcium signals may in itself influence embryonic patterning by regulating early embryonic cleavages. In ascidians, nemerteans and mouse, the egg cortex is polarized along the animal-vegetal axis and, in ascidians, this polarity amplifies after fertilization through actomyosin-driven cortical contractions (Sardet et al., 2002). Is the generation of repetitive  $\text{Ca}^{2+}$  waves from the vegetal cortical pacemaker a mechanism used to prime the vegetal pole region for later developmental events such as cleavage or gastrulation, which, in nemerteans and ascidians, takes place in the vegetal/dorsal pole of the embryo? Mouse embryos were long thought to have no significant polarity until the late cleavage stage, but recent marking experiments show that in fact, as in ascidians and nemerteans, although regulation can override this polarity, there is a relationship between the animal-vegetal axis, the sperm entry point and the developmental axes of pre- and post-implantation embryos (reviewed in Lu et al., 2001).

Finding out whether  $\text{Ca}^{2+}$  wave patterns play a role in later development will require studies that interfere with the normal spatio-temporal pattern of  $\text{Ca}^{2+}$  waves without perturbing mitosis and cleavage. The rather simple ascidian embryo, which displays two different meiotic  $\text{Ca}^{2+}$  wave pacemakers and develops into a swimming tadpole within a day, is particularly suited to studies of the relationship between meiotic  $\text{Ca}^{2+}$  waves and development (Fig. 2) (Dumollard and Sardet, 2001). It should be possible in the future to relate patterns of  $\text{Ca}^{2+}$  waves and phenotypic differences in embryos.

We thank Mark Larman and Karl Swann for their help with imaging calcium waves in mouse eggs injected with sperm extracts. We are also grateful to Christian Rouviere and Mohammed Khamla for technical assistance.

### References

- Albrieux, M., Moutin, M. J., Grunwald, D. and Villaz, M. (2000). Calmodulin and immunophilin are required as functional partners of a ryanodine receptor in ascidian oocytes at fertilization. *Dev. Biol.* **225**, 101-111.
- Albritton, N. L., Meyer, T. and Stryer, L. (1992). Range of messenger action of calcium ions and IP<sub>3</sub>. *Science* **258**, 1812-1815.
- Arnoult, C. and Villaz, M. (1994). Differential developmental fates of the two calcium currents in early embryos of the ascidian *Ciona intestinalis*. *J. Membr. Biol.* **137**, 127-135.
- Arnoult, C., Grunwald, D. and Villaz, M. (1996). Novel postfertilization inward  $\text{Ca}^{2+}$  current in ascidian eggs ensuring a calcium entry throughout meiosis. *Dev. Biol.* **174**, 322-334.
- Ayabe, T., Kopf, G. S. and Schultz, R. M. (1995). Regulation of mouse egg activation: presence of ryanodine receptors and effects of microinjected ryanodine and cyclic ADP ribose on uniseminated and inseminated eggs. *Development* **121**, 2233-2244.
- Berridge, M. J. (1997). Elementary and global aspects of calcium signalling. *J. Physiol.* **499**, 291-306.
- Boitier, E., Rea, R. and Duchen, M. (1999). Mitochondria exert a negative feedback on the propagation of intracellular  $\text{Ca}^{2+}$  waves in rat cortical astrocytes. *J. Cell Biol.* **145**, 795-808.
- Bootman, M. D., Lipp, P. and Berridge, M. J. (2001). The organisation and functions of local  $\text{Ca}^{2+}$  signals. *J. Cell Sci.* **114**, 2213-2222.
- Brind, S., Swann, K. and Carroll, J. (2000). Inositol 1,4,5-trisphosphate receptors are downregulated in mouse oocytes in response to sperm or adenophostin A but not to increases in intracellular  $\text{Ca}^{2+}$  or egg activation. *Dev. Biol.* **223**, 251-265.
- Bugrim, A. E., Zhabotinsky, A. M. and Epstein, I. R. (1997). Calcium waves in a model with a random spatially discrete distribution of  $\text{Ca}^{2+}$  release sites. *Biophys. J.* **73**, 2897-2906.
- Calarco, P. G. (1995). Polarization of mitochondria in the unfertilised mouse oocyte. *Dev. Genet.* **16**, 36-43.
- Carroll, J. (2000).  $\text{Na}^{+}$ - $\text{Ca}^{2+}$  exchange in mouse oocytes: modifications in the regulation of intracellular free  $\text{Ca}^{2+}$  during oocyte maturation. *J. Reprod. Fertil.* **118**, 337-342.
- Carroll, J. (2001). The initiation and regulation of  $\text{Ca}^{2+}$  signalling at fertilization in mammals. *Semin. Cell Dev. Biol.* **12**, 37-43.
- Chiba, K., Kado, R. T. and Jaffe, L. A. (1990). Development of calcium release mechanisms during starfish oocyte maturation. *Dev. Biol.* **140**, 300-306.
- Csutora, P., Su, Z., Kim, H. Y., Bugrim, A. and Cunningham, K. W. (1999). Calcium influx factor is synthesized by yeast and mammalian cells depleted of organellar calcium stores. *Proc. Natl. Acad. Sci. USA* **96**, 121-126.
- Deguchi, R., Osanai, K. and Morisawa, M. (1996). Extracellular  $\text{Ca}^{2+}$  entry and  $\text{Ca}^{2+}$  release from inositol 1,4,5-trisphosphate-sensitive stores function at fertilization in oocytes of the marine bivalve *Mytilus edulis*. *Development* **122**, 3651-3660.
- Deguchi, R., Shirakawa, H., Oda, S., Mohri, T. and Miyazaki, S. (2000). Spatiotemporal analysis of  $\text{Ca}^{2+}$  waves in relation to the sperm entry site and animal-vegetal axis during  $\text{Ca}^{2+}$  oscillations in the fertilised mouse eggs. *Dev. Biol.* **218**, 299-313.
- Duchen, M. (2000). Mitochondria and calcium: from cell signalling to cell death. *J. Physiol.* **529**, 57-68.
- Dumollard, R. and Sardet, C. (2001). Three different calcium wave pacemakers in ascidian eggs. *J. Cell Sci.* **114**, 2471-2481.
- Eisen, A. and Reynolds, G. T. (1985). Source and sinks for the calcium released during fertilization of single sea urchin eggs. *J. Cell Biol.* **100**, 1522-1527.
- Eckberg, W. R. and Miller, A. L. (1995). Propagated and non propagated calcium transients during egg activation in the annelid, Chaetopterus. *Dev. Biol.* **172**, 654-664.
- Fissore, R. A., Longo, F. J., Anderson, E., Parys, J. B. and Ducibella, T. (1999). Differential distribution of inositol trisphosphate receptor isoforms in mouse oocytes. *Biol. Reprod.* **60**, 49-57.
- Gillot, I., Ciapa, B., Payan, P. and Sardet, C. (1991). The calcium content of cortical granules and the loss of calcium from sea urchin eggs at fertilization. *Dev. Biol.* **146**, 396-405.
- Girard, J., Gillot, I., Renzis, G. D. and Payan, P. (1991). Calcium pools in sea urchin eggs: roles of endoplasmic reticulum mitochondria in relation to fertilization. *Cell. Calcium* **12**, 289-299.
- Halet, G., Tunwell, R., Balla, T., Swann, K. and Carroll, J. (2002). The dynamics of plasma membrane PtdIns(4,5)P<sub>2</sub> at fertilization of mouse eggs. *J. Cell Sci.* **115**, 2139-2149.
- Hirose, K., Kadowaki, S., Tanabe, M., Takeshima, H. and Iino, M. (1999). Spatiotemporal dynamics of inositol trisphosphate that underlies complex calcium mobilization patterns. *Science* **284**, 1527-1530.
- Ito, K., Miyashita, Y. and Kasai, H. (1999). Kinetic control of multiple forms of  $\text{Ca}^{2+}$  spikes by inositol trisphosphate in pancreatic acinar cells. *J. Cell Biol.* **146**, 405-413.
- Jacob, R. (1990). Calcium oscillations in electrically non-excitabile cells. *Biochem. Biophys. Acta* **1052**, 427-428.
- Jaconi, M., Pyle, J., Bortolon, R., Ou, J. and Clapham, D. (1997). Calcium release and influx colocalize to the endoplasmic reticulum. *Curr. Biol.* **7**, 599-602.
- Jaffe, L. F. (1991). The path of calcium in cytosolic calcium oscillations: a unifying hypothesis. *Proc. Natl. Acad. Sci. USA* **88**, 9883-9887.
- Jaffe, L. A. and Terasaki, M. (1993). Structural changes of the endoplasmic reticulum of sea urchin eggs during fertilization. *Dev. Biol.* **156**, 566-573.
- Jaffe, L. A., Giusti, A. F., Carroll, D. J. and Foltz, K. R. (2001).  $\text{Ca}^{2+}$

- signalling during fertilization of echinoderm eggs. *Semin. Cell. Dev. Biol.* **12**, 45-51.
- Jellerette, T., He, C. L., Wu, H., Parys, J. B. and Fissore, R. A.** (2000). Down-regulation of the inositol 1,4,5-trisphosphate receptor in mouse eggs following fertilization or parthenogenetic activation. *Dev. Biol.* **223**, 238-250.
- Jones, K.** (1998).  $Ca^{2+}$  oscillations in the activation of the egg and development of the embryo in mammals. *Int. J. Dev. Biol.* **42**, 1-10.
- Jones, K. and Nixon, V.** (2000). Sperm-induced  $Ca^{2+}$  oscillations in mouse oocytes and eggs can be mimicked by photolysis of caged inositol 1,4,5-trisphosphate: evidence to support a continuous low level production of inositol 1,4,5-trisphosphate during mammalian fertilization. *Dev. Biol.* **225**, 1-12.
- Kasai, H. and Petersen, O. H.** (1994). Spatial dynamics of second messengers: IP3 and cAMP as long-range and associative messengers. *Trends Neurosci.* **17**, 95-101.
- Kline, D., Mehlmann, L., Fox, C. and Terasaki, M.** (1999). The cortical endoplasmic reticulum (ER) of the mouse egg: localization of ER clusters in relation to the generation of repetitive calcium waves. *Dev. Biol.* **215**, 431-442.
- Kuthreiber, W. M., Gillot, I., Sardet, C. and Jaffe, L. F.** (1993). Net calcium and acid release at fertilization in eggs of sea urchins and ascidians. *Cell. Calcium* **14**, 73-86.
- Kyozuka, K., Deguchi, R., Mohri, T. and Miyazaki, S.** (1998). Injection of sperm extract mimics spatiotemporal dynamics of  $Ca^{2+}$  responses and progression of meiosis at fertilization of ascidian oocytes. *Development* **125**, 4099-4105.
- Leclerc, C., Guerrier, P. and Moreau, M.** (2000). Role of dihydropyridine-sensitive calcium channels in meiosis and fertilization in the bivalve molluscs *Ruditapes philippinarum* and *Crassostrea gigas*. *Biol. Cell* **92**, 285-299.
- Lee, M. G., Xu, X., Zheng, W., Diaz, J. and Wojcikiewicz, R. J.** (1997). Polarized expression of  $Ca^{2+}$  channels in pancreatic and salivary gland cells. *J. Biol. Chem.* **272**, 15765-15770.
- Liu, L., Hammar, K., Smith, P. J., Inoue, S. and Keefe, D. L.** (2001). Mitochondrial modulation of calcium signaling at the initiation of development. *Cell. Calcium* **30**, 423-433.
- Lu, C. C., Brennan, J. and Robertson, E. J.** (2001). From fertilization to gastrulation: axis formation in the mouse embryo. *Curr. Opin. Genet. Dev.* **11**, 384-392.
- Machaca, K. and Haun, S.** (2000). Store-operated calcium entry inactivates at the germinal vesicle breakdown stage of *Xenopus* meiosis. *J. Biol. Chem.* **275**, 38710-38715.
- Mak, D., McBride, S. and Foskett, J.** (1999). ATP regulation of type 1 inositol 1,4,5-trisphosphate receptor channel gating by allosteric tuning of  $Ca^{2+}$  activation. *J. Biol. Chem.* **274**, 22231-22237.
- Mak, D., McBride, S. and Foskett, J.** (2001). ATP regulation of recombinant type 3 inositol 1,4,5-trisphosphate receptor gating. *J. Gen. Physiol.* **117**, 447-456.
- Marchant, J. S. and Parker, I.** (2001). Role of elementary  $Ca^{2+}$  puff in generating repetitive  $Ca^{2+}$  oscillations. *EMBO J.* **20**, 65-76.
- Marchant, J. S., Callamaras, N. and Parker, I.** (1999). Initiation of  $InsP_3$ -mediated calcium waves in *Xenopus* oocytes. *EMBO J.* **18**, 5285-5299.
- McDougall, A. and Sardet, C.** (1995). Function and characteristics of repetitive calcium waves associated with meiosis. *Curr. Biol.* **5**, 318-328.
- McDougall, A., Shearer, J. and Whitaker, M.** (2000). The initiation and propagation of the fertilization wave in sea urchin eggs. *Biol. Cell* **92**, 205-214.
- McGuinness, O. M., Moreton, R. B., Johnson, M. H. and Berridge, M. J.** (1996). A direct measurement of increased divalent cation influx in fertilised mouse oocytes. *Development* **122**, 2199-2206.
- McPherson, S. M., McPherson, P. S., Mathews, L., Campbell, K. P. and Longo, F. J.** (1992). Cortical localization of a calcium release channel in sea urchin eggs. *J. Cell Biol.* **116**, 1111-1121.
- Mehlmann, L. M. and Kline, D.** (1994). Regulation of intracellular calcium in the mouse egg: calcium release in response to sperm or inositol trisphosphate is enhanced after meiotic maturation. *Biol. Reprod.* **51**, 1088-1098.
- Mehlmann, L. M., Mikoshiba, K. and Kline, D.** (1996). Redistribution and increase in cortical inositol 1,4,5-trisphosphate receptors after meiotic maturation of the mouse oocyte. *Dev. Biol.* **180**, 489-498.
- Mehlmann, L. M., Chattopadhyay, A., Carpenter, G. and Jaffe, L. A.** (2001). Evidence that phospholipase C from the sperm is not responsible for initiating  $Ca^{2+}$  release at fertilization in mouse eggs. *Dev. Biol.* **236**, 492-501.
- Meyer, T. and Stryer, L.** (1988). Molecular model for receptor-stimulated calcium spiking. *Proc. Natl. Acad. Sci. USA* **85**, 5051-5055.
- Miyazaki, S., Shirakawa, H., Nakada, K. and Honda, Y.** (1993). Essential role of the inositol trisphosphate receptor/ $Ca^{2+}$  release channel in  $Ca^{2+}$  waves and  $Ca^{2+}$  oscillations at fertilization of mammalian eggs. *Dev. Biol.* **158**, 62-78.
- Nash, M. S., Young, K. W., Willars, G. B., Challiss, R. A. J. and Nahorski, S. R.** (2001). Single-cell imaging and graded  $Ins(1,4,5)P_3$  production following G-protein-coupled-receptor activation. *Biochem. J.* **356**, 137-142.
- Nixon, V. L., McDougall, A. and Jones, K. T.** (2000). Calcium oscillations and the cell cycle at fertilization of mammalian and ascidian eggs. *Biol. Cell* **92**, 187-196.
- Oda, S., Deguchi, R., Mohri, T., Shikano, T., Nakanishi, S. and Miyazaki, S.** (1999). Spatiotemporal dynamics of the  $[Ca^{2+}]_i$  rise induced by microinjection of sperm extracts into mouse eggs: preferential induction of a  $Ca^{2+}$  wave from the cortex mediated by the inositol 1,4,5-trisphosphate receptor. *Dev. Biol.* **209**, 172-185.
- Parrington, J., Lai, F. A. and Swann, K.** (2000). The soluble mammalian sperm factor protein that triggers  $Ca^{2+}$  oscillations in eggs: evidence for expression of mRNA(s) coding for sperm factor protein(s) in spermatogenic cells. *Biol. Cell* **92**, 267-275.
- Petersen, O. H., Burdakov, D. and Tepikin, A. V.** (1999). Polarity in intracellular calcium signalling. *BioEssays* **21**, 851-860.
- Putney, J. W., Jr, Broad, L. M., Braun, F. J., Lievreumont, J. P. and Bird, G. S.** (2001). Mechanisms of capacitative calcium entry. *J. Cell Sci.* **114**, 2223-2239.
- Rice, A., Parrington, J., Jones, K. T. and Swann, K.** (2000). Mammalian sperm contain a  $Ca^{2+}$ -sensitive phospholipase C activity that can generate  $InsP(3)$  from  $PIP(2)$  associated with intracellular organelles. *Dev. Biol.* **228**, 125-135.
- Rizzuto, R., Bernardi, P. and Pozzan, T.** (2000). Mitochondria as all-round players of the calcium game. *J. Physiol.* **529**, 37-47.
- Rooney, T. A., Sass, E. J. and Thomas, A. P.** (1990). Agonist-induced cytosolic calcium oscillations originate from a specific locus in single hepatocytes. *J. Biol. Chem.* **265**, 10792-10796.
- Runft, L. L. and Jaffe, L. A.** (2000). Sperm extract injection into ascidian eggs signals  $Ca^{2+}$  release by the same pathway as fertilization. *Development* **127**, 3227-3236.
- Runft, L. L., Watras, J. and Jaffe, L. A.** (1999). Calcium release at fertilization of *Xenopus* eggs requires type I  $IP(3)$  receptors, but not SH2 domain-mediated activation of PLCgamma or G(q)-mediated activation of PLCbeta. *Dev. Biol.* **214**, 399-411.
- Runft, L. L., Jaffe, L. A. and Mehlmann, L. M.** (2002). Egg activation at fertilization: where it all begins. *Dev. Biol.* **245**, 237-254.
- Rutter, G. A. and Rizzuto, R.** (2000). Regulation of mitochondrial metabolism by ER  $Ca^{2+}$  release: an intimate connection. *Trends Biochem. Sci.* **25**, 215-221.
- Sardet, C., Roegiers, F., Dumollard, R., Rouviere, C. and McDougall, A.** (1998). Calcium waves and oscillations in eggs. *Biophys. Chem.* **72**, 131-140.
- Sardet, C., Prodon, F., Dumollard, R., Chang, P. and Chenevert, J.** (2002). Structure and function of the egg cortex. *Dev. Biol.* **241**, 1-23.
- Saunders, C. M., Larman, M. G., Parrington, J., Cox, L. J., Royse, J., Blayney, L. M., Swann, K. and Lai, F. A.** (2002).  $PLC\zeta$ : a sperm-specific trigger of  $Ca^{2+}$  oscillations in eggs and embryo development. *Development* **129**, 3533-3544.
- Shiraishi, K., Okada, A., Shirakawa, H., Nakanishi, S., Mikoshiba, K. and Miyazaki, S.** (1995). Developmental changes in the distribution of the endoplasmic reticulum and inositol 1,4,5-trisphosphate receptors and the spatial pattern of  $Ca^{2+}$  release during maturation of hamster oocytes. *Dev. Biol.* **170**, 594-606.
- Simpson, P. B., Mehotra, S., Lange, G. D. and Russell, J. T.** (1997). High density distribution of endoplasmic reticulum proteins and mitochondria at specialized  $Ca^{2+}$  release sites in oligodendrocyte processes. *J. Biol. Chem.* **272**, 22654-22661.
- Speknijder, J.** (1992). The repetitive calcium waves in the fertilised ascidian egg are initiated near the vegetal pole by a cortical pacemaker. *Dev. Biol.* **153**, 259-271.
- Speknijder, J., Sardet, C. and Jaffe, L.** (1990a). The activation wave of calcium in the ascidian egg and its role in ooplasmic segregation. *J. Cell Biol.* **110**, 1589-1598.
- Speknijder, J., Sardet, C. and Jaffe, L.** (1990b). Periodic calcium waves cross ascidian eggs after fertilization. *Dev. Biol.* **142**, 246-249.

- Speksnijder, J., Terasaki, M., Hage, W., Jaffe, L. and Sardet, C.** (1993). Polarity and reorganisation of the endoplasmic reticulum during fertilization and ooplasmic segregation in the ascidian egg. *J. Cell Biol.* **120**, 1337-1346.
- Stricker, S.** (1999). Comparative biology of calcium signaling during fertilization and egg activation in animals. *Dev. Biol.* **211**, 157-176.
- Stricker, S. A., Silva, R. and Smythe, T.** (1998). Calcium and endoplasmic reticulum dynamics during oocyte maturation and fertilization in the marine worm *Cerebratulus lacteus*. *Dev. Biol.* **203**, 305-322.
- Swann, K. and Parrington, J.** (1999). Mechanism of Ca<sup>2+</sup> release at fertilization in mammals. *J. Exp. Zool.* **285**, 267-275.
- Taylor, C. W., Genazzani, A. A. and Morris, S. A.** (1999). Expression of inotistol trisphosphate receptors. *Cell. Calcium* **26**, 237-251.
- Terasaki, M., Jaffe, L. A., Hunnicutt, G. R. and Hammer, J. A., 3rd** (1996). Structural change of the endoplasmic reticulum during fertilization: evidence for loss of membrane continuity using the green fluorescent protein. *Dev. Biol.* **179**, 320-328.
- Terasaki, M., Runft, L. L. and Hand, A. R.** (2001). Changes in organisation of the endoplasmic reticulum during *Xenopus* oocyte maturation and activation. *Mol. Biol. Cell* **12**, 1103-1116.
- Thomas, D., Lipp, P., Tovey, S. C., Berridge, M. J. and Li, W.** (1999). Microscopic properties of elementary calcium release sites in non-excitabile cells. *Curr. Biol.* **18**, 8-15.
- Van Blerkom, J., Davis, P., Mathwig, V. and Alexander, S.** (2002). Domains of high-polarized and low-polarized mitochondria may occur in mouse and human oocytes and early embryos. *Hum. Reprod.* **17**, 393-406.